

A CYCLIC EXTENSION OF THE EARTHQUAKE FLOW

FRANCESCO BONSANTE, GABRIELE MONDELLO, AND JEAN-MARC SCHLENKER

ABSTRACT. Let \mathcal{T} be Teichmüller space of a closed surface of genus at least 2. For any point $c \in \mathcal{T}$, we describe an action of the circle on $\mathcal{T} \times \mathcal{T}$, which limits to the earthquake flow when one of the parameters goes to a measured lamination in the Thurston boundary of \mathcal{T} . This circle action shares some of the main properties of the earthquake flow, for instance it satisfies an extension of Thurston's Earthquake Theorem and it has a complex extension which is analogous and limits to complex earthquakes. Moreover, a related circle action on $\mathcal{T} \times \mathcal{T}$ extends to the product of two copies of the universal Teichmüller space.

CONTENTS

1. Introduction	2
1.1. Earthquakes on hyperbolic surfaces	2
1.2. Harmonic maps and the landslide flow	3
1.3. Minimal Lagrangian maps between hyperbolic surfaces	4
1.4. The landslide action on $\mathcal{T} \times \mathcal{T}$	4
1.5. Relations to AdS geometry	4
1.6. The center of circles in Teichmüller space	5
1.7. Obtaining R from L	5
1.8. The earthquake flow as a limit	5
1.9. An extension of the Earthquake Theorem	6
1.10. A complex extension	6
1.11. Landslide on the universal Teichmüller space	6
1.12. Content of the paper	7
Acknowledgment	7
2. Minimal lagrangian maps and AdS geometry	7
2.1. Notations	7
2.2. Hyperbolic ends	7
2.3. The duality between hyperbolic and de Sitter ends	8
2.4. Globally hyperbolic AdS manifolds	8
2.5. Minimal Lagrangian maps between hyperbolic surfaces	9
3. Definition of the cyclic flow	9
3.1. The Levi-Civita connection of h_θ	10
3.2. A cyclic property	10
3.3. Proof of Theorem 1.8	11
3.4. Proof of Theorem 1.10	11
3.5. Centers	12
4. An extension of Thurston's Earthquake Theorem	12
4.1. Constant curvature foliations in AdS geometry	12
4.2. Proof of Theorem 1.14	12
5. The complex cyclic map	13
5.1. Main statements	13
5.2. General formulas	14
5.3. The variation field of SGr'	16
5.4. The comparison of the cocycles	18
5.5. Parameterization by the disk	19

Date: June 2011 (v1).

F.B. is partially supported by the A.N.R. through project Geodycos .

J.-M. S. was partially supported by the A.N.R. through projects ETTT, ANR-09-BLAN-0116-01, and GeomEinstein, ANR-09-BLAN-0116-01.

6. The earthquake flow as a limit	20
6.1. Convergence on the imaginary axis	21
6.2. Convergence on the real axis	22
6.3. Asymptotic behavior of the measures $\text{tr}(b_n)\omega_{h_n}$	27
6.4. Proof of Theorems 1.12 and 6.1	29
6.5. Convergence of the distances	30
7. Behavior of the centers	33
8. The landslide flow on the universal Teichmüller space	36
8.1. Minimal Lagrangian maps and the universal Teichmüller space	36
8.2. An extension of L to the universal Teichmüller space	36
9. Applications, extensions, and questions	37
9.1. Constant curvature surfaces in globally hyperbolic AdS manifolds	37
9.2. Holomorphic disks in Teichmüller space	37
9.3. Other questions	37
References	38

1. INTRODUCTION

In all the paper we consider a closed, oriented surface S of genus at least 2. We denote by \mathcal{T}_S , or simply by \mathcal{T} , Teichmüller space of S , and by \mathcal{ML}_S , or simply by \mathcal{ML} , the space of measured laminations on S .

1.1. Earthquakes on hyperbolic surfaces. Given a measured lamination $\lambda \in \mathcal{ML}_S$, we denote by E_λ the *left earthquake* along λ on S . E_λ is a real-analytic map from \mathcal{T}_S to \mathcal{T}_S , see [32, 11, 19]. Recall that, in the simplest case where λ is supported on the simple closed curve γ with mass a , if $h \in \mathcal{T}_S$ is a hyperbolic metric on S , $E_\lambda(h)$ is obtained by cutting (S, h) open along the minimizing geodesic homotopic to γ , rotating the left-hand side of γ by a , and gluing back.

We consider here the *earthquake flow*, which can be defined as a map:

$$\begin{aligned} E : \mathbb{R} \times \mathcal{T} \times \mathcal{ML} &\rightarrow \mathcal{T} \times \mathcal{ML} \\ (t, h, \lambda) &\mapsto (E_{t\lambda}(h), \lambda) \end{aligned}$$

We call E_t the corresponding map from $\mathcal{T} \times \mathcal{ML}$ to $\mathcal{T} \times \mathcal{ML}$, and will also use the notation $E_\lambda(h) := E_1(h, \lambda)$. Earthquakes have a number of interesting properties, of which we can single the following.

- (1) The earthquake flow defined above is indeed a flow: for all $s, t \in \mathbb{R}$, $E_s \circ E_t = E_{s+t}$.
- (2) Thurston's Earthquake Theorem (see [11]): for any $h, h' \in \mathcal{T}$, there is a unique $\lambda \in \mathcal{ML}$ such that $E_\lambda(h) = h'$.
- (3) For fixed $\lambda \in \mathcal{ML}_S$ and $h \in \mathcal{T}_S$, the map

$$\begin{aligned} \mathbb{R} &\rightarrow \mathcal{T}_S \\ t &\mapsto E_{-t\lambda}(h) \end{aligned}$$

extends to a holomorphic map on a simply connected domain in \mathbb{C} containing all complex numbers with non-negative imaginary part, see [19]. This defines the notion of “complex earthquake”.

- (4) When considered on imaginary numbers, complex earthquakes correspond to *grafting* maps. The conformal grafting map $gr : \mathbb{R}_{\geq 0} \times \mathcal{T} \times \mathcal{ML} \rightarrow \mathcal{T}$ is actually obtained by composing the projective grafting map

$$Gr : \mathbb{R}_{\geq 0} \times \mathcal{T} \times \mathcal{ML} \rightarrow \mathcal{P},$$

where \mathcal{P} is the space of complex projective structures on S , with the forgetful map $\mathcal{P} \rightarrow \mathcal{T}$ sending a \mathbb{CP}^1 -structure to the underlying complex structure.

- (5) Thurston (see [14]) proved that, for all $s > 0$, Gr_s is a homeomorphism from $\mathcal{T} \times \mathcal{ML}$ to \mathcal{P} .

We introduce a flow on Teichmüller space which in a way extends the earthquake flow, and which shares the properties described above. The corresponding deformations are “smoother” than earthquakes, but earthquakes are limits in a natural sense. This motivates the term “landslide” that we use here. This deformation depends not on a measured lamination but rather on a hyperbolic metric $h^* \in \mathcal{T}$ and it determines a flow:

$$\begin{aligned} L : \mathcal{T} \times \mathcal{T} \times S^1 &\rightarrow \mathcal{T} \times \mathcal{T} \\ (h, h^*, e^{i\theta}) &\mapsto L_{e^{i\theta}, h^*}(h). \end{aligned}$$

We denote by $L_{e^{i\theta}} : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T} \times \mathcal{T}$ the corresponding map seen as depending on the parameter $e^{i\theta}$.

We will also use the notation L^1 for the composition of L with the projection on the first factor, so that L^1 is a map from $S^1 \times \mathcal{T} \times \mathcal{T}$ to \mathcal{T} . $L_{e^{i\theta}}^1$ will denote the same map, considered as depending on the parameter $e^{i\theta} \in S^1$, so it is a map from $\mathcal{T} \times \mathcal{T}$ to \mathcal{T} . Thus $L_{e^{i\theta}}^1$ is the analog of the earthquake map $E : \mathcal{T} \times \mathcal{ML} \rightarrow \mathcal{T}$.

When the metrics h and h^* are fixed, we can consider the image of the map $S^1 \ni e^{i\theta} \mapsto L_{e^{i\theta}}^1(h, h^*) \in \mathcal{T}$ as a circle in Teichmüller space in which h and h^* are antipodal points. When h^* converges (projectively) to a measured lamination λ at Thurston boundary of Teichmüller space, such a circle converges to the earthquake line $E_{t\lambda}(h)$. A more precise statement, Theorem 1.12, can be found below.

This “landslide flow” shares the main properties of the earthquake flow recalled above:

- (1) L is a flow on $\mathcal{T} \times \mathcal{T}$ – depending on the definition, checking this can be non trivial, see Theorem 1.8.
- (2) We prove an analog of Thurston’s Earthquake Theorem, see Theorem 1.14.
- (3) For fixed $h, h^* \in \mathcal{T}$, the map $L_\bullet(h, h^*) : S^1 \rightarrow \mathcal{T}$ extends to a holomorphic map from the closed unit disk $\overline{\Delta}$ to \mathcal{T} , see Theorem 5.1. This defines the “complex landslide” which are analogs of the “complex earthquakes”.
- (4) For $r \in (0, 1)$, the complex landslide L_r corresponds to what we call here “smooth grafting”, which is analog to grafting in our context and we denote by $sgr_r : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$ the map defined as $L_r : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T} \times \mathcal{T}$ followed by projection on the first factor. It is obtained by composing a map $SGr : (0, 1) \times \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{P}$ with the natural projection from \mathcal{P} to \mathcal{T} .
- (5) For all $r \in (0, 1)$, the map $SGr(r, \bullet, \bullet) : (0, 1) \times \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{P}$ is a homeomorphism.

Our notations mean that we parameterize the complex landslides using the unit disk in \mathbb{C} rather than the upper half-plane as is customary for complex earthquakes. This notation is clearly equivalent but using the disk appears more natural in the context of the landslides considered here.

Considered as a circle action on $\mathcal{T} \times \mathcal{T}$, the flow L extends to a circle action on the universal Teichmüller space, see Section 8.

1.2. Harmonic maps and the landslide flow. Consider two hyperbolic metrics c and h on S . A map $f : (S, c) \rightarrow (S, h)$ is said to be *harmonic* if it is a critical point of the energy E . The energy considered here is:

$$E(f) = \frac{1}{2} \int_S \|df\|^2 \omega_c$$

where ω_c is the area element of (S, c) . Although it is not immediately apparent in this definition, this notion of harmonicity is conformally invariant on the domain, so that we can regard c as a conformal structure on S rather than a metric.

Theorem 1.1 (Sampson [24], Schoen and Yau [29]). *Let c be a conformal class on S , and let $h \in \mathcal{T}$ be a hyperbolic metric. There is a unique harmonic map $f : (S, c) \rightarrow (S, h)$ isotopic to the identity. Moreover, f is a diffeomorphism.*

Consider a C^1 map $f : (S, h^\#) \rightarrow (S, h)$, where $h^\#$ is a metric in the conformal class of c . The Hopf differential $\Phi(f)$ of f is a quadratic differential that measures the traceless part of the pull-back of h by f and it is defined by the formula

$$f^*h = eh^\# + \Phi + \overline{\Phi},$$

where $e = \frac{1}{2} \text{tr}_{h^\#}(f^*h)$. If f is harmonic, then Φ is holomorphic. For $f \in C^2$, also the converse holds. It follows from its definition that $\Phi(f)$ is invariant under conformal changes of the metric $h^\#$ on S .

Conversely, given a holomorphic quadratic differential Φ on (S, c) , there exists a unique hyperbolic metric h on S such that the identity map $(S, c) \rightarrow (S, h)$ is harmonic with Hopf differential Φ , see [24, 33].

This leads to the definition of a flow on \mathcal{T} depending on a “center” $c \in \mathcal{T}$.

Definition 1.2. Let $c, h \in \mathcal{T}$ and let $e^{i\theta} \in S^1$. We define $R_{c, e^{i\theta}}(h)$ as the (unique) hyperbolic metric h' on S such that, if $f : (S, c) \rightarrow (S, h)$ and $f' : (S, c) \rightarrow (S, h')$ are the harmonic maps isotopic to the identity, then

$$\Phi(f') = e^{i\theta} \Phi(f).$$

This simple definition is strongly related to the flow L mentioned above, but the relation is not obvious (see Corollary 1.11), and using directly the definition of R given here is not convenient. For this reason we give below a different definition of L , which is more geometric, less directly accessible, but leads to straightforward arguments.

There is another, superficially similar flow on Teichmüller space, the elliptic flow defined by one of us (Mon-dello), see [21]. There are only limited similarities between the two flows, as should be clear from the sequel.

1.3. Minimal Lagrangian maps between hyperbolic surfaces. The constructions considered here depend strongly on the notion of minimal Lagrangian map between hyperbolic surfaces. Recall that, given two hyperbolic metrics h and h^* on S , a diffeomorphism $m : (S, h) \rightarrow (S, h^*)$ is *minimal Lagrangian* if:

- it is area-preserving and orientation-preserving,
- its graph is a minimal surface in $(S \times S, h \oplus h^*)$.

Theorem 1.3 (Schoen [30], Labourie [16]). *Let h, h^* be two hyperbolic metrics on S . There exists a unique minimal Lagrangian diffeomorphism $m : (S, h) \rightarrow (S, h^*)$ isotopic to the identity.*

Minimal Lagrangian maps actually have a description in terms of hyperbolic surfaces only, as follows (see e.g. [16]).

Proposition 1.4. *If $m : (S, h) \rightarrow (S, h^*)$ is minimal Lagrangian, then $m^*h^* = h(b\bullet, b\bullet)$, where $b : TS \rightarrow TS$:*

- (1) *is self-adjoint for h ,*
- (2) *has determinant 1,*
- (3) *satisfies the Codazzi equation: $d^\nabla b = 0$, where ∇ is the Levi-Civita connection of h .*

Conversely, if $m : S \rightarrow S$ is a diffeomorphism satisfying those properties, then it is minimal Lagrangian.

Corollary 1.5. *Let h, h^* be two hyperbolic metrics on S . There exists a unique bundle morphism $b : TS \rightarrow TS$ which is self-adjoint for h , of determinant equal to 1 everywhere, satisfies Codazzi equation $d^\nabla b = 0$, where ∇ is the Levi-Civita connection of h , and such that $h(b\bullet, b\bullet)$ is isotopic to h^* .*

A consequence of this proposition is that for any $\tau, \tau^* \in \mathcal{T}$, we can realize τ and τ^* as a pair of hyperbolic metrics h and h^* (not considered up to isotopies) so that $h^* = h(b\bullet, b\bullet)$, where b is self-adjoint for h , of determinant 1, and satisfies Codazzi equation $d^\nabla b = 0$. A pair of metrics with this property will be called a *normalized representative* of (τ, τ^*) . Clearly a normalized representative of (τ, τ^*) is uniquely determined up to isotopies acting diagonally on both h and on h^* .

By abuse of notation, we will sometimes denote by (h, h^*) both a couple of normalized hyperbolic metrics and its correspondent point in $\mathcal{T} \times \mathcal{T}$.

1.4. The landslide action on $\mathcal{T} \times \mathcal{T}$. We now introduce the action L of S^1 on $\mathcal{T} \times \mathcal{T}$. We will see below that it is strongly related to the map R introduced above.

Definition 1.6. Let h, h^* be two hyperbolic metrics on S , and let $\theta \in \mathbb{R}$. We consider the bundle morphism $b : TS \rightarrow TS$ given by Corollary 1.5, and set

$$(1) \quad \beta_\theta := \cos(\theta/2)E + \sin(\theta/2)Jb ,$$

where $E : TS \rightarrow TS$ is the identity map and J is the complex structure of h on S . We then call

$$L_{e^{i\theta}}(h, h^*) := (h(\beta_\theta\bullet, \beta_\theta\bullet), h(\beta_{\theta+\pi}\bullet, \beta_{\theta+\pi}\bullet)) .$$

Notice that by construction, $L_1(h, h^*) = (h, h^*)$, while $L_{-1}(h, h^*) = (h^*, h)$.

Proposition 1.7. *For all $\theta \in \mathbb{R}$, $h(\beta_\theta\bullet, \beta_\theta\bullet)$ is a hyperbolic metric on S .*

Theorem 1.8. *Let h, h^* be two hyperbolic metrics on S , let $\theta, \theta' \in \mathbb{R}$. Then*

$$L_{e^{i\theta'}}(L_{e^{i\theta}}(h, h^*)) = L_{e^{i(\theta'+\theta)}}(h, h^*) .$$

*In other terms, L defines an action of S^1 on $\mathcal{T} \times \mathcal{T}$. We call L the **landslide flow**, or **landslide action** on $\mathcal{T} \times \mathcal{T}$.*

The proofs of Proposition 1.7 and of Theorem 1.8 are in Section 3.3.

1.5. Relations to AdS geometry. We briefly recall some properties of globally hyperbolic anti-de Sitter manifolds. More details can be found e.g. in [20, 1].

The anti-de Sitter space is a Lorentz analog of hyperbolic 3-space, it can be defined as the quadric:

$$\text{AdS}^3 = \{x \in \mathbb{R}^{2,2} \mid \langle x, x \rangle = -1\} ,$$

where $\mathbb{R}^{2,2} = (\mathbb{R}^4, -dx_0^2 - dx_1^2 + dx_2^2 + dx_3^2)$. It is a complete Lorentz manifold of constant curvature -1 with fundamental group isomorphic to \mathbb{Z} .

A manifold N with an AdS metric – a Lorentz metric locally modeled on AdS^3 – is *maximal globally hyperbolic* (MGH) if:

- N contains a closed space-like surface F ,
- any inextendible time-like curve in N intersects F exactly once,

- N is maximal for inclusion, under these properties.

Mess [20, 1] proved that, if N is (GH) and $\bar{\phi} : S \rightarrow N$ is an embedding onto a closed space-like surface F , then N is the quotient of a convex domain Ω in AdS^3 by an action of the fundamental group of S .

A key feature of AdS^3 is that the identity component of its isometry group is isomorphic to $\text{SL}_2(\mathbb{R}) \times \text{SL}_2(\mathbb{R})/\mathbb{Z}_2$, which is the double cover of $\text{PSL}_2(\mathbb{R}) \times \text{PSL}_2(\mathbb{R})$. As a consequence, the action of $\pi_1(S)$ on Ω decomposes as (ρ_l, ρ_r) , where ρ_l and ρ_r are morphisms from $\pi_1(S)$ to $\text{PSL}_2(\mathbb{R})$. It was proved in [20] that these morphisms have maximal Euler number, so that they correspond to points in the Teichmüller space of S . Maximal globally hyperbolic AdS spaces are uniquely determined by these left and right representations, see [20, 1].

Lemma 1.9. *Let h, h^* be a pair of normalized metrics, let $\theta \in (0, \pi)$. There exists a unique equivariant embedding (ϕ, ρ) of \tilde{S} in AdS^3 with induced metric $\cos^2(\theta/2)h$ and third fundamental form $\sin^2(\theta/2)h^*$. Moreover, ρ is the holonomy representation of a globally hyperbolic AdS manifold N : the first factor in $L_{e^{i\theta}}(h, h^*)$ is the left representation of N and the first factor in $L_{e^{-i\theta}}(h, h^*)$ is the right representation of N .*

The proof is in Section 2.4.

1.6. The center of circles in Teichmüller space. Let $h, h^* \in \mathcal{T}$. For each $\theta \in \mathbb{R}$, let $(h_\theta, h_\theta^*) = L_{e^{i\theta}}(h, h^*)$. According to Theorem 1.3, there is a unique minimal Lagrangian diffeomorphism $m_\theta : (S, h_\theta) \rightarrow (S, h_\theta^*)$ isotopic to the identity. We can then consider on S the conformal structure c_θ of the metric $h_\theta + m_\theta^* h_\theta^*$. We call c_θ the *center* of (h_θ, h_θ^*) . This conformal class of metrics has some interesting properties, proved in Section 3.4.

Theorem 1.10. (1) m_θ is the identity – that is, the identity is minimal Lagrangian between (S, h_θ) and (S, h_θ^*) .
 (2) c_θ does not depend on θ , it is equal to a fixed conformal class c .
 (3) Let $f_\theta : (S, c) \rightarrow (S, h_\theta)$ and $f_\theta^* : (S, c) \rightarrow (S, h_\theta^*)$ be the unique harmonic maps isotopic to the identity. Then f_θ and f_θ^* have opposite Hopf differentials.
 (4) For any $\theta \in \mathbb{R}$,

$$\Phi(f_\theta) = e^{i\theta} \Phi(f_0) .$$

1.7. Obtaining R from L . As a consequence of Theorem 1.10, we find a simple relation between the map R defined earlier in terms of Hopf differential, and the map L .

Corollary 1.11. *Let (h, h^*) be a couple of normalized metrics and let c be the conformal class of $h + h^*$. For any $e^{i\theta} \in S^1$, we have*

$$L_{e^{i\theta}}(h, h^*) = (R_{c, \theta}(h), R_{c, \theta + \pi}(h^*)) .$$

1.8. The earthquake flow as a limit.

Theorem 1.12. *Let $h \in \mathcal{T}$, let $(h_n^*)_{n \in \mathbb{N}}$ be a sequence of hyperbolic metrics and let $\lambda \neq 0$ be a measured lamination. Consider a sequence $(\theta_n)_{n \in \mathbb{N}}$ of positive real numbers such that $\lim_{n \rightarrow \infty} \theta_n \ell_{h_n^*} = \iota(\lambda, \bullet)$ in the sense of convergence of the length spectra of simple closed curves. Then*

$$\lim_{n \rightarrow +\infty} h_n^1 = E_{\lambda/2}(h) \quad \lim_{n \rightarrow +\infty} \theta_n \ell_{h_n^2} = \iota(\lambda, \bullet)$$

where $(h_n^1, h_n^2) := L_{e^{i\theta_n}}(h, h_n^*)$.

At first sight it would appear more natural to take the limit where the sequence of centers (c_n) converges projectively to λ . However the statement obtained by replacing L by R and h_n^* by c_n in Theorem 1.12 turns out to be false, as proved – in one example – in Section 7, see Corollary 7.3.

The heuristic argument motivating Theorem 1.12 involves the convergence of constant Gauss curvature surfaces to a pleated surface in AdS^3 . However, writing a proof based on these ideas turns out to be more difficult than it appears. A key technical statement is that minimal Lagrangian maps have a close proximity to Thurston compactification of Teichmüller space: minimal Lagrangian maps isotopic to the identity provide “the” correct normalization to understand the convergence of a sequence of hyperbolic metrics to a projective measured lamination in Thurston boundary of \mathcal{T} .

Theorem 1.13. *Let h be a hyperbolic metric on S , and let $(h_n^*)_{n \in \mathbb{N}}$ be a sequence of hyperbolic metrics such that $\theta_n \ell_{h_n^*} \rightarrow \iota(\lambda, \bullet)$, where λ is a measured geodesic lamination, the θ_n are positive numbers, and the convergence is in the sense of the length spectrum. For each n , let $m_n : (S, h) \rightarrow (S, h_n^*)$ be the minimal Lagrangian diffeomorphism isotopic to the identity. Then, for every smooth arc α in S that meets the h -geodesic representative of λ transversely and with endpoints not in the support of (the h -geodesic representative of) λ , the length for $\theta_n^2 m_n^*(h_n^*)$ of the geodesic segment homotopic to α (with fixed endpoints) converges to the intersection between α and λ .*

The proof of this theorem involves the convergence of smooth surfaces to a pleated limit, but in the hyperbolic, rather than the anti-de Sitter, context.

1.9. An extension of the Earthquake Theorem. We can now state an extension to the landslide flow L of Thurston's Earthquake Theorem (see [11]). Recall that this theorem states that, given two hyperbolic metrics h and h' on a surface, there is a unique measured lamination λ such that the left earthquake along λ sends h to h' .

Theorem 1.14. *Let $h, h' \in \mathcal{T}$ and let $e^{i\theta} \in S^1 \setminus \{0\}$. There is a unique $h^* \in \mathcal{T}$ such that $L_{e^{i\theta}}^1(h, h^*) = h'$.*

We give in section 4.2 a simple proof based on a recent result of Barbot, Béguin and Zeghib [3] on the existence and uniqueness of constant Gauss curvature foliations in globally hyperbolic AdS manifolds.

As an easy consequence, a similar statement holds also for the flow R .

1.10. A complex extension. The earthquake flow has an extension as a map $E : \overline{\mathbb{H}} \times \mathcal{T} \times \mathcal{ML} \rightarrow \mathcal{T}$, where $\overline{\mathbb{H}}$ is the set of complex numbers with nonnegative imaginary part. This map has the property that, for any $h \in \mathcal{T}$ and any $\lambda \in \mathcal{ML}$, the map $z \mapsto E(z, h, \lambda)$ is holomorphic, see [19]. It can be defined in terms of grafting, or (for small λ) in terms of pleated surfaces in hyperbolic 3-space.

In Section 5 we prove that the landslide map L defined above has a similar holomorphic extension where the parameter $e^{i\theta}$ is replaced by a complex number ζ in the closed unit disk. This defines many holomorphic disks in Teichmüller space, see Theorem 5.1. Similarly to what happens for complex earthquakes, this construction factors through the space of complex projective structures on S for $\zeta \neq 0$, and the complex cyclic flow provides punctured holomorphic disks in this space. This factorization however does not extend for $\zeta = 0$.

The complex landslide map limits to complex earthquakes just as the “real” landslide flow limits to the earthquake flow, see Theorem 6.1.

We hope at some point in the future to give another proof of the holomorphicity of this complex landslide map, based on a geometric argument taking place in the complexification of \mathbb{H}^3 . This line of argument should also provide a straightforward and geometric way to understand why complex earthquakes are holomorphic disks.

1.11. Landslide on the universal Teichmüller space. Recall that a homeomorphism of the circle is *quasi-symmetric* if and only if it is the boundary value of a quasi-conformal diffeomorphism of the disk.

Definition 1.15. The universal Teichmüller space \mathcal{T}_U is the quotient of the group \mathcal{QS} of quasi-symmetric homeomorphisms of the circle by left composition by projective transformations.

The universal Teichmüller space contains embedded copies of the Teichmüller space of all closed surfaces. Indeed, consider a closed surface S of genus at least 2, a fixed hyperbolic metric $h^\#$ on S , and its holonomy representation $\rho^\# : \pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbb{R})$. Given another hyperbolic metric h on S and its holonomy representation $\rho : \pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbb{R})$, there is a quasiconformal map $\tilde{f} : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ conjugating $\rho^\#$ and ρ . Moreover, the boundary value $\partial\tilde{f} : \partial_\infty\mathbb{H}^2 \rightarrow \partial_\infty\mathbb{H}^2$ is uniquely determined by $\rho^\#$ and ρ , and the map sending h to $\partial\tilde{f}$ is an embedding of \mathcal{T}_S in \mathcal{T}_U , see e.g. [10].

Let $\psi : S^1 \rightarrow S^1$ be a quasi-symmetric homeomorphism. There is (see [7]) a unique minimal Lagrangian quasiconformal diffeomorphism $m : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ with $\partial m = \psi$. As for closed surfaces, there is then a unique bundle morphism $b : T\mathbb{H}^2 \rightarrow T\mathbb{H}^2$ such that

- b is self-adjoint,
- it satisfies the Codazzi equation $d^\nabla b = 0$,
- $m^*g = g(b\bullet, b\bullet)$, where g is the hyperbolic metric on \mathbb{H}^2 .

For every $\theta \in \mathbb{R}$ we then consider $\beta_\theta := \cos(\theta/2)E + \sin(\theta/2)b$, where E is the identity and $g_\theta := g(\beta_\theta\bullet, \beta_\theta\bullet)$.

Lemma 1.16. *g_θ is a complete hyperbolic metric on \mathbb{H}^2 . The identity map between (\mathbb{H}^2, g) and (\mathbb{H}^2, g_θ) is quasiconformal (and minimal Lagrangian), and its extension $\psi_\theta : S^1 \rightarrow S^1$ to the boundary of \mathbb{H}^2 is quasi-symmetric, so that it defines a point in \mathcal{QS} .*

In Section 8 we show how to use this fact to construct an extension of L to a non-trivial circle action \mathcal{L} on $\mathcal{T}_U \times \mathcal{T}_U$ (see Theorem 8.5).

1.12. Content of the paper. In Section 2 we present the background material, concerning in particular minimal Lagrangian diffeomorphisms between hyperbolic surfaces and globally hyperbolic AdS manifolds. In Section 3 we define the landslide flow and prove that it is indeed a flow (Theorem 1.8) as well as Theorem 1.10. In Section 4 we give the proof of the extension to the landslide flow of Thurston's Earthquake Theorem (Theorem 1.14). Then in Section 5 we construct the complex landslide map, actually as a map from $\mathcal{T} \times \mathcal{T} \times \dot{\bar{\Delta}}$ to \mathcal{P} , where $\dot{\bar{\Delta}}$ is the pointed closed unit disk in \mathbb{C} , and we prove that it is holomorphic and that it extends over $\bar{\Delta}$ as a map to \mathcal{T} . Section 6 considers the limit when the parameter h^* converges projectively to a measured lamination at Thurston boundary of Teichmüller space, and contains the proof of Theorem 1.12 as well as its complex extension, Theorem 6.1. In Section 7, on the other hand, we show that the situation is not as simple for the “center” c determined by a fixed metric h and a sequence h_n^* going to a point at infinity in Thurston compactification of \mathcal{T} : the limit of the corresponding sequence of centers does not depend only on the limit of (h_n^*) . Section 8 deals with the circle action on the universal Teichmüller space, while Section 9 contains some remarks and open questions.

Acknowledgment. We are grateful to the Institute of Mathematical Sciences of the National University of Singapore, where most of the results presented here were obtained. The second named author would like to thank Mike Wolf for helpful clarifications on his paper [34].

2. MINIMAL LAGRANGIAN MAPS AND ADS GEOMETRY

We present in this section some background material used in the paper.

2.1. Notations. In all the paper we consider a closed, oriented surface S of genus at least 2.

We consider AdS^3 , as well as all AdS manifolds, as oriented and time-oriented. All the embeddings of S that we consider will implicitly be considered as time-oriented, that is, the oriented normal to the image is future-oriented. Moreover, the convex embeddings will always be considered to be positively convex, that is, the oriented normal is future-directed and pointing towards the convex side. We recall that it is possible to identify the isometries of AdS^3 with double cover of $\text{PSL}_2(\mathbb{R}) \times \text{PSL}_2(\mathbb{R})$ in such a way that, if S is a positively convex pleated surface in AdS^3 , bent along λ and with first fundamental form h , then the first (resp. second) factor corresponds to the holonomy of the hyperbolic surface obtained from h performing a left (resp. right) earthquake along λ .

2.2. Hyperbolic ends. The 3-dimensional hyperbolic space can be defined as a quadric in the 4-dimensional Minkowski space $\mathbb{R}^{1,3} = (\mathbb{R}^4, -dx_0^2 + dx_1^2 + dx_2^2 + dx_3^2)$, with the induced metric.

$$\mathbb{H}^3 = \{x \in \mathbb{R}^{1,3} \mid \langle x, x \rangle = -1 \text{ and } x_0 > 0\}.$$

It is a simply connected, complete manifold with constant curvature -1 .

A quasifuchsian hyperbolic manifold is a 3-dimensional manifold locally isometric to \mathbb{H}^3 , homeomorphic to $S \times \mathbb{R}$, which contains a non-empty compact convex subset.

Such a quasifuchsian manifold M contains a smallest non-empty convex subset $C(M)$ called its convex core. M is *Fuchsian* if $C(M)$ is a totally geodesic surface, otherwise the boundary of $C(M)$ is the disjoint union of two pleated surfaces.

Each connected component of the complement of $C(M)$ in M is an instance of a *hyperbolic end*: a hyperbolic manifold homeomorphic to $S \times \mathbb{R}_{>0}$, complete on one side and bounded by a locally concave pleated surface on the other. There is a one-to-one correspondence between hyperbolic ends homeomorphic to $S \times \mathbb{R}_{>0}$ and complex projective structures on S , which associates to a hyperbolic end the natural complex projective structure on its boundary at infinity, see e.g. [14].

Labourie [16] proved that any hyperbolic end has a unique foliation by convex, constant curvature surfaces. The curvature varies monotonically from -1 close to the pleated surface boundary, to 0 close to the boundary at infinity.

Given an oriented surface Σ in a hyperbolic end M (or in \mathbb{H}^3) we will usually denote by I its induced metric, and by B its shape operator, considered as a bundle morphism from $T\Sigma$ to $T\Sigma$. It is defined by $BX = \nabla_X \nu$, where ν is the oriented unit normal to Σ and ∇ is the Levi-Civita connection of M . We will also denote by $E : TS \rightarrow TS$ the identity.

Definition 2.1. Let Σ be a convex surface embedded in a hyperbolic end M with embedding data (I_Σ, B_Σ) . The *grafted metric* on Σ is $I_\Sigma^\# = I_\Sigma((E + B_\Sigma)\bullet, (E + B_\Sigma)\bullet)$.

A basic and well-known property of this metric $I_\Sigma^\#$ is that the hyperbolic Gauss map – sending a point $x \in \Sigma$ to the endpoint at infinity of the geodesic ray starting at x orthogonal to Σ – is a conformal map between $(\Sigma, I_\Sigma^\#)$ and $\partial_\infty M$ with its conformal structure. More details will be found in Section 6.

2.3. The duality between hyperbolic and de Sitter ends. The 3-dimensional de Sitter space can be defined, as the hyperbolic space, as a quadric in the 4-dimensional Minkowski space, with the induced metric.

$$dS^3 = \{x \in \mathbb{R}^{1,3} \mid \langle x, x \rangle = 1\}.$$

There is a one-to-one correspondence between points in dS^3 and oriented totally geodesic planes in \mathbb{H}^3 , see e.g. [23, 27]. Given an oriented surface $S \subset \mathbb{H}^3$, its dual is the set S^* of points of dS^3 corresponding to oriented planes tangent to S in \mathbb{H}^3 . If S is smooth and locally strictly convex, then S^* is also smooth and locally strictly convex.

Consider a quasifuchsian hyperbolic manifold M , and let E be one of the ends of M , that is, one of the connected components of the complement of $C(M)$ in M . The universal cover of M is identified with \mathbb{H}^3 , and the universal cover \tilde{E} of E is then identified with a connected component of the complement of the convex hull of the limit set Λ of $\pi_1(M)$ in $\partial_\infty \mathbb{H}^3$. Let \tilde{E}^* be the set of points of dS^3 corresponding to oriented planes in \mathbb{H}^3 contained in \tilde{E} . Then (see [20]) $\pi_1(M)$ acts properly discontinuously on \tilde{E}^* , and the quotient is a de Sitter *domain of dependence*, that is, a globally hyperbolic maximal de Sitter manifold (see below for the definition in the AdS case).

This construction actually extends (see [20]) to a hyperbolic end E which is not necessarily one of the ends of a quasifuchsian manifold. In this manner, any hyperbolic end E has a “dual” de Sitter domain of dependence E^* , and conversely.

One feature of this duality which will be used below is that if S is a surface in E with constant curvature K , then there is a dual surface S^* in E^* . (It is the quotient by $\pi_1(M)$ of the surface in dS^3 dual to the universal cover of S in \mathbb{H}^3 .) The curvature of S^* is then also constant, and equal to $K/(K+1)$. In this manner a foliation of E by constant curvature surfaces gives rise to a dual foliation of E^* by constant curvature surfaces (see [3] for more details).

2.4. Globally hyperbolic AdS manifolds. The definition of AdS^3 and of globally hyperbolic AdS manifolds are recalled in the introduction. There are many similarities between quasifuchsian hyperbolic manifolds and globally hyperbolic AdS manifolds, some of which – being used in the arguments below – are recalled here.

Let N be a globally hyperbolic AdS 3-manifold, and let F be a closed, space-like surface in N for which the induced metric has negative sectional curvature (or, equivalently, the determinant of the second fundamental form of F is everywhere larger than -1). Let I and ν be the induced metric on this surface and a unit normal vector. Let J be the complex structure induced by ν on F : namely $J(v) = \nu \times v$ where \times is the vector product on $T\text{AdS}^3$. Finally, let $B = \nabla \nu$ be the shape operator of F , where ∇ is the Levi-Civita connection of AdS^3 . We consider the Riemannian metrics on F

$$h_l = I((E + JB)\bullet, (E + JB)\bullet), h_r = I((E - JB)\bullet, (E - JB)\bullet).$$

Then h_l and h_r are two smooth hyperbolic metrics on F (see [12]). This can be used when F is a maximal or constant mean curvature surface in N , but also when F is a constant Gauss curvature surface.

Remark 2.2. Notice that, even if J and B depend on the choice of a normal vector, JB and the metrics h_l and h_r are independent of it.

According to our orientation and time-orientation of AdS^3 , the holonomy of the metric h_l is equal to the first factor of the holonomy of N and the holonomy of h_r is equal to the second factor of the holonomy of N , see [12]. This last observation can be used to prove Lemma 1.9.

Proof of Lemma 1.9. Let $b : TS \rightarrow TS$ be the positive h -self-adjoint operator given by Corollary 1.5 such that $h^* = h(b\bullet, b\bullet)$. Since h and h^* are normalized metrics we know that b is a solution of Codazzi equation and $\det b = 1$.

We consider now the pair

$$I_\theta = \cos^2(\theta/2)h \quad B_\theta = \tan(\theta/2)b.$$

Clearly, B_θ is a solution of Codazzi equation. Moreover, since I_θ is a metric of constant curvature $K = -\frac{1}{\cos^2(\theta/2)}$, we easily get that

$$K_\theta = -1 - \det B_\theta$$

so (I_θ, B_θ) is also a solution of Gauss equation for spacelike AdS surfaces (see [12]).

This implies that there is an equivariant map

$$\phi : \tilde{S} \rightarrow \tilde{F} \subset \text{AdS}^3$$

whose embedding data are \tilde{I}_θ and \tilde{B}_θ . The map ϕ is unique up to isometries of AdS^3 . We will also require that

- the normal field $\tilde{\nu}$ that induces the right orientation on \tilde{S} points toward the convex side
- $\tilde{\nu}$ is a future-directed vector field.

□

Lemma 1.9 and [12] imply that $L_{e^{i\theta}}(h, h^*)$ is a couple of hyperbolic metrics. For convenience of the reader, we will give a simple proof of this fact in Section 3.1.

Globally hyperbolic AdS manifolds have a unique foliation by constant mean curvature surfaces, see [2].

A globally hyperbolic AdS manifold N contains a smallest non-empty convex subset $C(N)$, called its convex core: N is called *Fuchsian* if $C(N)$ is a totally geodesic surface; otherwise, the boundary of $C(N)$ is the disjoint union of two pleated locally convex surfaces in N , so that its induced metric is hyperbolic and its pleating is described by a measured lamination (see [20]). The complement of $C(N)$ in N has two connected components, one future convex, the other past convex. Barbot, Béguin and Zeghib [3] proved that $N \setminus C(N)$ has a unique foliation by convex, constant Gauss curvature surfaces. The curvature is monotonic along the foliation, and varies from -1 in the neighborhood of the convex core to $-\infty$ near the initial/final singularity.

2.5. Minimal Lagrangian maps between hyperbolic surfaces. The definition of minimal Lagrangian diffeomorphisms has been recalled in the introduction. Remark that the definition directly shows that the inverse of a minimal Lagrangian diffeomorphism is also minimal Lagrangian.

Let us mention here that they occur in several distinct geometric contexts, and the interplay between the different occurrences is used below, in particular in Section 4.

- If S is a surface of constant curvature K in a constant curvature, Riemannian or Lorentzian 3-manifold M , then the third fundamental form $\mathbb{I}\!\!\mathbb{I}$ of S also has constant curvature K^* , where K^* depends on K , on the ambient curvature, and on whether the ambient space is Riemannian or Lorentzian (for instance, if $M = \mathbb{H}^3$, then $K^* = K/(K+1)$). If both K and K^* are negative, then $|K|I$ and $|K^*|\mathbb{I}\!\!\mathbb{I}$ are hyperbolic metrics, and the identity, considered as a map from $(S, |K|I)$ to $(S, |K^*|\mathbb{I}\!\!\mathbb{I})$, is minimal Lagrangian.
- If M is an “almost Fuchsian” manifold, that is, M is a quasifuchsian hyperbolic 3-manifolds containing a closed, embedded minimal surface S with principal curvatures everywhere in $(-1, 1)$, then S is the unique closed minimal surface in M . The “hyperbolic Gauss maps” send S to each connected component of the boundary at infinity of M , and both maps are diffeomorphisms. Composing these maps one finds a diffeomorphism between one component of $\partial_\infty M$ and the other. This diffeomorphism is minimal Lagrangian if each boundary component is endowed with the (unique) hyperbolic metric in its conformal class. (See e.g. [12] for details and proofs.)
- Similarly, if N is a globally hyperbolic AdS manifold, then it contains a unique closed, space-like maximal surface F . Consider the metrics h_l and h_r defined above on F . Then h_l and h_r are the left and right hyperbolic metrics of N , respectively, and moreover the identity between (F, h_l) and (F, h_r) is minimal Lagrangian (see [12] for details).

It is the first of these occurrences which will play the largest role here.

Minimal Lagrangian maps between hyperbolic surfaces are intimately related to harmonic maps: let $m : (S, h) \rightarrow (S', h')$ be a minimal Lagrangian diffeomorphism between two hyperbolic surfaces and consider the conformal structure c on S of the metric $h + m^*h'$. Then

- the identity is harmonic between (S, c) and (S, h) ,
- m is harmonic between (S, c) and (S', h') ,
- those two harmonic maps have opposite Hopf differentials.

The converse is also true. Details can be found e.g. in [30].

3. DEFINITION OF THE CYCLIC FLOW

In this section we consider two fixed normalized hyperbolic metrics h, h^* on S , and call b the bundle morphism given by Corollary 1.5. Let β_θ be the family of operators defined in (1).

Definition 3.1. Given $\theta \in \mathbb{R}$, we call

$$h_\theta = h(\beta_{\theta\bullet}, \beta_{\theta\bullet}) .$$

Comparing with definition 1.6, we have $L_{e^{i\theta}}(h, h^*) = (h_\theta, h_{\theta+\pi})$.

3.1. The Levi-Civita connection of h_θ .

Lemma 3.2. *For all $\theta \in \mathbb{R}$, $d^\nabla \beta_\theta = 0$, where ∇ is the Levi-Civita connection of h .*

Proof. Let u, v be two vector fields on S . Note first that β_θ satisfies Codazzi equation:

$$\begin{aligned} (d^\nabla \beta_\theta)(u, v) &= \nabla_u(\beta_\theta v) - \nabla_v(\beta_\theta u) - \beta_\theta([u, v]) \\ &= \cos(\theta/2)(\nabla_u v - \nabla_v u - [u, v]) + \sin(\theta/2)(\nabla_u(Jbv) - \nabla_v(Jbu) - Jb[u, v]) \\ &= \sin(\theta/2)J(\nabla_u(bv) - \nabla_v(bu) - b[u, v]) \\ &= 0, \end{aligned}$$

where the last equality follows from the fact that b satisfies Codazzi equation. \square

Lemma 3.3. *The Levi-Civita connection ∇^θ of h_θ is given by*

$$\nabla_u^\theta v = \beta_{-\theta} \nabla_u(\beta_\theta v).$$

Proof. Consider the connection ∇^θ defined in the statement of the lemma. It is sufficient to prove that it is compatible with h_θ and torsion-free.

Let u, v, w be three vector fields on S . Then

$$\begin{aligned} u \cdot h_\theta(v, w) &= u \cdot h(\beta_\theta v, \beta_\theta w) \\ &= h(\nabla_u(\beta_\theta v), \beta_\theta w) + h(\beta_\theta v, \nabla_u(\beta_\theta w)) \\ &= h_\theta(\nabla_u^\theta v, w) + h_\theta(v, \nabla_u^\theta w), \end{aligned}$$

and therefore ∇^θ is compatible with h_θ .

We can now compute the torsion of ∇^θ on u, v .

$$\begin{aligned} \nabla_u^\theta v - \nabla_v^\theta u - [u, v] &= \beta_{-\theta}(\nabla_u(\beta_\theta v) - \nabla_v(\beta_\theta u) - \beta_\theta[u, v]) \\ &= \beta_{-\theta}(d^\nabla \beta_\theta)(u, v) \\ &= 0. \end{aligned}$$

Thus ∇^θ is torsion-free and compatible with h_θ , and so it is the Levi-Civita connection of h_θ . \square

The computation of the connection of h_θ allows to prove easily that h_θ is a hyperbolic metric.

Proof of Proposition 1.7. Let (e_1, e_2) be an orthonormal frame for h on S , with connection 1-form ω with respect to the Levi-Civita connection ∇ of h . Let $(e'_1, e'_2) = (\beta_\theta^{-1} e_1, \beta_\theta^{-1} e_2)$, it is an orthonormal moving frame for h_θ . Since the Levi-Civita connection ∇^θ of h_θ is given by Lemma 3.3, a direct computation shows that the connection 1-form of (e'_1, e'_2) for ∇^θ is also equal to ω . Since the identity map between (S, h) and (S, h_θ) is area-preserving, it follows that h and h_θ have the same curvature, so that h_θ is also hyperbolic. \square

3.2. A cyclic property. Our next goal is to prove Theorem 1.8 and Theorem 1.10. The proofs are in the next sections, after some preliminary lemmas.

Lemma 3.4. *Let $\theta, \theta' \in \mathbb{R}$. Then $\beta_\theta \circ \beta_{\theta'} = \beta_{\theta+\theta'}$.*

Proof. By definition we have:

$$\begin{aligned} \beta_\theta \circ \beta_{\theta'} &= (\cos(\theta/2)E + \sin(\theta/2)Jb) \circ (\cos(\theta'/2)E + \sin(\theta'/2)Jb) \\ &= \cos(\theta/2)\cos(\theta'/2)E + \sin(\theta/2)\sin(\theta'/2)JbJb + (\cos(\theta/2)\sin(\theta'/2) + \sin(\theta/2)\cos(\theta'/2))Jb. \end{aligned}$$

But $JbJb = -E$ because b is self-adjoint and of determinant 1. It follows that

$$\beta_\theta \circ \beta_{\theta'} = \cos((\theta + \theta')/2)E + \sin((\theta + \theta')/2)Jb = \beta_{\theta+\theta'}.$$

\square

For instance, since $Jb = \beta_\pi$, it follows that $Jb \circ \beta_\theta = \beta_\theta \circ Jb = \beta_{\theta+\pi}$.

Lemma 3.5. *Let $b_\theta : TS \rightarrow TS$ be the bundle morphism associated to h_θ and h_θ^* by Corollary 1.5. Then*

$$b_\theta = \beta_{-\theta} \circ b \circ \beta_\theta.$$

Proof. Let $b_\theta = \beta_{-\theta} \circ b \circ \beta_\theta$. This lemma will follow if we prove the following facts:

- (1) $h_\theta(b_\theta \bullet, b_\theta \bullet)$ is isotopic to h_θ^* (we will actually prove that it is equal to h_θ^*).
- (2) b_θ has determinant 1.
- (3) b_θ is self-adjoint for h_θ .

$$(4) \quad d^{\nabla^\theta} \beta_\theta = 0.$$

For the first point, note that $Jb = \beta_\pi$, and therefore:

$$\begin{aligned} h_\theta(b_\theta \bullet, b_\theta \bullet) &= h(b\beta_\theta \bullet, b\beta_\theta \bullet) = h(Jb\beta_\theta \bullet, Jb\beta_\theta \bullet) \\ &= h(\beta_\pi \circ \beta_\theta \bullet, \beta_\pi \circ \beta_\theta \bullet) = h(\beta_{\pi+\theta} \bullet, \beta_{\pi+\theta} \bullet) \\ &= h_\theta^* . \end{aligned}$$

The second point is clear since $\det(b_\theta) = \det(\beta_{-\theta}) \det(b) \det(\beta_\theta) = 1$.

For point (3) let u, v be two vector fields on S , then

$$\begin{aligned} h_\theta(b_\theta u, v) &= h(\beta_\theta b_\theta u, \beta_\theta v) = h(b\beta_\theta u, \beta_\theta v) \\ &= h(\beta_\theta u, b\beta_\theta v) = h_\theta(u, b_\theta v) . \end{aligned}$$

For point (4), let again be u, v be two vector fields on S . Then, using the expression of ∇^θ in Lemma 3.3, we have:

$$\begin{aligned} (d^{\nabla^\theta} b_\theta)(u, v) &= \nabla_u^\theta(b_\theta v) - \nabla_v^\theta(b_\theta u) - b_\theta[u, v] \\ &= \beta_{-\theta} \nabla_u(\beta_\theta b_\theta v) - \beta_{-\theta} \nabla_v(\beta_\theta b_\theta u) - b_\theta[u, v] \\ &= \beta_{-\theta}(d^\nabla \beta_{\theta+\pi})(u, v) \end{aligned}$$

so that $(d^{\nabla^\theta} b_\theta)(u, v) = 0$ by Lemma 3.2. This completes the proof. \square

Remark 3.6. Lemma 3.5 shows that h_θ and $h_{\theta+\pi}$ are normalized metrics.

3.3. Proof of Theorem 1.8. Let $\theta, \theta' \in \mathbb{R}$. By definition of $L_{\theta'}$,

$$L_{\theta'}(L_\theta(h, h^*)) = (h_\theta(\bar{\beta}_{\theta'} \bullet, \bar{\beta}_{\theta'} \bullet), (h_{\theta+\pi}(\bar{\beta}_{\theta'} \bullet, \bar{\beta}_{\theta'} \bullet))) ,$$

where

$$\bar{\beta}_{\theta'} = \cos(\theta'/2)E + \sin(\theta'/2)J_\theta b_\theta ,$$

where J_θ is the complex structure of h_θ . Clearly, $J_\theta = \beta_{-\theta} \circ J \circ \beta_\theta$, so that

$$\begin{aligned} \bar{\beta}_{\theta'} &= \beta_{-\theta} \circ (\cos(\theta'/2)E + \sin(\theta'/2)Jb) \circ \beta_\theta \\ &= \beta_{-\theta} \circ \beta_{\theta'} \circ \beta_\theta \\ &= \beta_{\theta'} . \end{aligned}$$

We now see that

$$\begin{aligned} L_{\theta'}(L_\theta(h, h^*)) &= (h_\theta(\beta_{\theta'} \bullet, \beta_{\theta'} \bullet), (h_{\theta+\pi}(\beta_{\theta'} \bullet, \beta_{\theta'} \bullet))) \\ &= (h(\beta_\theta \circ \beta_{\theta'} \bullet, \beta_\theta \circ \beta_{\theta'} \bullet), h(\beta_{\theta+\pi} \circ \beta_{\theta'} \bullet, \beta_{\theta+\pi} \circ \beta_{\theta'} \bullet)) \\ &= (h(\beta_{\theta'+\theta} \bullet, \beta_{\theta'+\theta} \bullet), h^*(\beta_{\theta'+\theta} \bullet, \beta_{\theta'+\theta} \bullet)) \\ &= L_{\theta'+\theta}(h, h^*) . \end{aligned}$$

This proves Theorem 1.8.

3.4. Proof of Theorem 1.10. The fact that h_θ and $h_{\theta+\pi}$ are normalized metrics follows from Lemma 3.5.

We compute the expression of c_θ .

$$\begin{aligned} c_\theta &= h_\theta + h_{\theta+\pi} \\ &= h((\cos(\theta/2)E + \sin(\theta/2)b)\bullet, (\cos(\theta/2)E + \sin(\theta/2)b)\bullet) + \\ &\quad + h((-\sin(\theta/2)E + \cos(\theta/2)b)\bullet, (-\sin(\theta/2)E + \cos(\theta/2)b)\bullet) \\ &= (\cos^2(\theta/2) + \sin^2(\theta/2))h + (\cos^2(\theta/2) + \sin^2(\theta/2))h(b\bullet, b\bullet) \\ &= c , \end{aligned}$$

so c_θ is indeed independent of θ .

The fact that the identity $(S, c) \rightarrow (S, h_\theta)$ is harmonic follows from the last paragraph of Subsection 2.5.

For point (4) recall that b_θ has determinant 1 and that $h_\theta^* = h_\theta(b_\theta \bullet, b_\theta \bullet)$. A simple computation then shows that $h_\theta - h_\theta^*$ is traceless with respect to $h_\theta + h_\theta^*$. The definition of Hopf differential (see Section 1.2) therefore shows that, if Φ_θ is the Hopf differential of the identity from (S, c) to (S, h_θ) , then

$$\frac{h_\theta - h_\theta^*}{2} = 2\text{Re}(\Phi_\theta) .$$

Therefore

$$\begin{aligned}
4\operatorname{Re}(\Phi_\theta) &= h((\cos(\theta/2)E + \sin(\theta/2)b)\bullet, (\cos(\theta/2)E + \sin(\theta/2)b)\bullet) - \\
&\quad - h((-\sin(\theta/2)E + \cos(\theta/2)b)\bullet, (-\sin(\theta/2)E + \cos(\theta/2)b)\bullet) \\
&= (\cos^2(\theta/2) - \sin^2(\theta/2))(h - h^*) + 4\sin(\theta/2)\cos(\theta/2)h(b\bullet, \bullet) \\
&= \cos(\theta)(h - h^*) + 2\sin(\theta)h(b\bullet, \bullet).
\end{aligned}$$

So $\operatorname{Re}(\Phi_\theta) = \operatorname{Re}(e^{i\theta}\Phi)$, where

$$(2) \quad \Phi = \frac{h - h^* + 2ih(b\bullet, \bullet)}{4} = \frac{h((E - b^2)\bullet, \bullet) + 2ih(b\bullet, \bullet)}{4}$$

is the Hopf differential of the identity from (S, c) to (S, h) .

3.5. Centers. We conclude this section by some remarks on the respective behavior of h, h^* and c .

Remark 3.7. c is uniquely determined by $h, h^* \in \mathcal{T}$. Conversely, h^* is uniquely determined by h and c .

Proof. Given h and h^* , we have seen that there is a unique minimal Lagrangian diffeomorphism $m : (S, h) \rightarrow (S, h^*)$ isotopic to the identity. By definition, c is the conformal class of $h + m^*h^*$.

For the second point, let $f : (S, c) \rightarrow (S, h)$ be the unique harmonic map, given by Theorem 1.1, and let Φ be its Hopf differential. There is then a unique harmonic map f^* from (S, c) to a hyperbolic surface (S, h^*) with Hopf differential equal to $-\Phi$. The content of Section 2.5 then indicates that $f^* \circ f^{-1}$ is minimal Lagrangian, so that c is the center of (h, h^*) . \square

4. AN EXTENSION OF THURSTON'S EARTHQUAKE THEOREM

In this section we recall a recent result of [3] on constant curvature foliations of GH AdS manifolds, and use it to prove Theorem 1.14.

4.1. Constant curvature foliations in AdS geometry. We recall here one of the main result of [3]. Let N be a MGH AdS 3-dimensional manifold, let $C(N)$ be its convex core.

Theorem 4.1 (Barbot, Béguin, Zeghib [3]). *The complement of $C(N)$ in N is foliated by surfaces of constant (Gauss) curvature. Moreover, for any $k \in (-\infty, -1)$, there exists a unique future-convex (resp. past-convex) surface of constant curvature k in N , and it is a leaf of the foliation.*

4.2. Proof of Theorem 1.14. We first translate Theorem 4.1 in terms of the landslide flow, using Lemma 1.9.

Corollary 4.2. *Choose $(\rho_l, \rho_r) \in \mathcal{T}$ and $\alpha \in (0, \pi)$. There exists a unique $(h, h^*) \in \mathcal{T} \times \mathcal{T}$ such that*

$$L_{e^{i\alpha}}^1(h, h^*) = \rho_l \text{ and } L_{e^{-i\alpha}}^1(h, h^*) = \rho_r.$$

Proof. It is a direct consequence of Theorem 4.1. Given ρ_l and ρ_r there is a unique GHMC AdS manifold $N \cong S \times \mathbb{R}$ of which they are the left and right representations, respectively. N contains a unique past-convex surface F with constant curvature $-1/\cos^2(\alpha/2)$, which comes with a diffeomorphism $\tilde{\phi} : S \rightarrow F$ (well-defined up to isotopy). We call I and III the induced metric and third fundamental form on S , respectively. Then III has constant curvature $-1/\sin^2(\alpha/2)$. We then set $h = (1/\cos^2(\alpha/2))I$, $h^* = (1/\sin^2(\alpha/2))\text{III}$, so that h and h^* are normalized hyperbolic metrics on S (see [12]). Lemma 1.9 then shows that $\rho_l = L_{e^{i\alpha}}^1(h, h^*)$, while $\rho_r = L_{e^{-i\alpha}}^1(h, h^*)$.

Conversely, given $h, h^* \in \mathcal{T}$ such that $\rho_l = L_{e^{i\alpha}}^1(h, h^*)$, while $\rho_r = L_{e^{-i\alpha}}^1(h, h^*)$, we can consider the unique equivariant embedding $\phi : \tilde{S} \rightarrow \operatorname{AdS}^3$ onto a past-convex surface \tilde{F} , with induced metric $\cos^2(\alpha/2)h$ and third fundamental form $\sin^2(\alpha/2)h^*$. Then \tilde{F} is the lift to AdS^3 of a past-convex surface F in a GHMC AdS manifold N , and the left and right representations of N are ρ_l and ρ_r by Lemma 1.9. This shows the uniqueness. \square

Proof of Theorem 1.14. Apply Corollary 4.2 with $\rho_r = h$, $\rho_l = h'$, and with $\alpha = \theta/2$. It shows there exists a unique $h_0 \in \mathcal{T}$ and a unique $h_0^* \in \mathcal{T}$ such that

$$(3) \quad h = L_{e^{-i\alpha}}^1(h_0, h_0^*), \quad h' = L_{e^{i\alpha}}^1(h_0, h_0^*).$$

As a consequence, putting $h^* = L_{e^{-i\alpha}}^2(h_0, h_0^*)$ we have $L_{e^{i\theta}}^1(h, h^*) = h'$.

Conversely given $h^* \in \mathcal{T}$ such that $L_{e^{i\theta}}^1(h, h^*) = h'$, letting $(h_0, h_0^*) = L_{e^{i\theta/2}}(h, h^*)$ we easily see that (3) is verified. The uniqueness in Corollary 4.2 therefore implies the uniqueness here. \square

5. THE COMPLEX CYCLIC MAP

This section describes a natural extension of the cyclic flow from a real to a complex parameter. This is analogous to the complex earthquake introduced by McMullen [19]. We will actually show in the next section that the “complex cyclic flow” introduced here limits, in a suitable sense, to the complex earthquake.

5.1. Main statements. Let \mathcal{P} be the space of complex projective surface on S . The space \mathcal{P} is naturally a complex manifold of real dimension $12g - 12$. Moreover the natural map $\mathcal{P} \rightarrow \mathcal{T}$ that associates to a complex projective surface the underlying complex structure is holomorphic. A projective structure is *Fuchsian* if its universal covering is projectively equivalent to a round disk in \mathbb{CP}^1 .

Let \mathbb{H} be the upper half-plane in \mathbb{C} . We define a map

$$P' : \mathbb{H} \times \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{P}$$

where

- For every fixed h, h^* the map $z \rightarrow P_z(h, h^*)$ is holomorphic.
- For t real, $P'_t(h, h^*)$ is the Fuchsian projective surface corresponding to $L_{e^{-it}}^1(h, h^*)$.

Notice that here z lives in the upper half-plane, while, in the introduction, the flow usually depended on a complex parameter ζ in the unit disk. Both parameterizations are quite useful here. Taking ζ in the unit disk is natural when dealing with the landslide flow, while taking z in the upper half-space is natural when thinking of complex earthquakes as a limit (since complex earthquakes are usually parameterized by the upper half-space). Until Section 5.4 we consider the parameterization by z in the upper half-space, while in Section 5.5 we will make the connection to the parameterization by ζ in the unit disk. We use a prime to denote the various maps when z is in the upper half-plane, this explains the notation P' above.

The construction of the map P' is the analog of the construction of the complex earthquake due to McMullen [19]. The first point is to define the analog of the grafting.

Given two normalized hyperbolic metrics h and h^* on \mathcal{T} , let b be the operator defined in Subsection 1.3. Given a positive number $s > 0$, we consider the metric $I_s = \cosh^2(s/2)h$ and the operator $B_s = -\tanh(s/2)b$. It is easy to see that (I_s, B_s) satisfies the Gauss-Codazzi equation for immersed surfaces in \mathbb{H}^3 , that is

$$d^\nabla B_s = 0, \quad K_s = -1 + \det B_s,$$

where ∇ is the Levi-Civita connection for I_s (which is equal to the Levi-Civita connection for h) and K_s is the curvature of I_s (which is constant and equal to $-1/\cosh^2(s/2)$).

As a consequence there exists a convex equivariant immersion

$$(4) \quad \sigma_s : \tilde{S} \rightarrow \mathbb{H}^3$$

whose first fundamental form is the pull-back \tilde{I}_s of I_s and whose shape operator is the pull-back \tilde{B}_s of B_s . This map σ_s is uniquely determined up to elements of $\text{PSL}_2(\mathbb{C})$, once we state that the orientation on \tilde{S} at $\sigma_s(\tilde{p})$ coincides with the orientation induced by the normal vector $\tilde{\nu}_s(\tilde{p})$ pointing towards the concave part (this is the reason why the sign of B_s is negative).

Given $\tilde{p} \in \tilde{S}$, let $dev_s(\tilde{p}) \in S_\infty^2 = \mathbb{CP}^1$ be the endpoint of the geodesic ray starting from $\sigma_s(\tilde{p})$ with velocity $\tilde{\nu}_s(\tilde{p})$. The map

$$dev_s : \tilde{S} \rightarrow \mathbb{CP}^1$$

is a developing map for a complex projective structure $SGr'_s(h, h^*)$ on S . (The notation SGr is used to keep in mind the analogy to the grafting map Gr).

Notice that, if $h = h^*$, then $b = E$ and $\sigma_s = d_s \circ \sigma_0$, where $d_s : \sigma_0(\tilde{S}) = \mathbb{H}^2 \rightarrow \mathbb{H}^3$ is the map associating to $\sigma_{s_0}(\tilde{p})$ the end-point of the geodesic segment of length s starting from $\sigma_{s_0}(\tilde{p})$ with velocity $\tilde{\nu}_{s_0}(\tilde{p})$. So, in this case, $SGr'_s(h, h)$ is the Fuchsian projective structure associated to h .

Finally, given a complex number $z = t + is$ with $s \geq 0$, we define

$$P'_z(h, h^*) = SGr'_s(L'_{-t})(h, h^*)$$

where $L'_{-t}(h, h^*) := L_{e^{-it}}(h, h^*)$.

Most of the remaining part of this section is devoted to proving the following theorem.

Theorem 5.1. *The map*

$$z \mapsto P'_z(h, h^*) \in \mathcal{P}$$

is holomorphic.

Composing P' with the forgetful map from \mathcal{P} to \mathcal{T} , we obtain for each z in the upper half-plane a map

$$C'_z : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$$

sending $(h, h^*) \in \mathcal{T} \times \mathcal{T}$ to the complex structure underlying the complex projective structure $P'_z(h, h^*)$.

Corollary 5.2. *The map $z \rightarrow C'_z(h, h^*)$ is holomorphic in the upper half-plane.*

This clearly follows from Theorem 5.1 since the forgetful map from \mathcal{P} to \mathcal{T} is holomorphic.

To prove Theorem 5.1, we will show that the holonomy ρ_z of P'_z , holomorphically depends on z . In fact the derivatives

$$\xi_z = \frac{\partial \rho_z}{\partial t}, \quad \eta_z = \frac{\partial \rho_z}{\partial s}$$

are $\mathfrak{sl}_2(\mathbb{C})$ -valued cocycles in $H^1(\pi_1(S), Ad \circ \rho_z)$ and we will show that

$$(5) \quad \eta_z = i\xi_z.$$

Let us remark that, since $P'_{z+t}(h, h^*) = P'_z(L_{e^{-it}}(h, h^*))$ for any z in the upper plane and t real, it is sufficient to check (5) at imaginary points $z_0 = is_0$.

To compute the cocycles we consider the family of convex immersions

$$\sigma_s, \tau_t : \tilde{S} \rightarrow \mathbb{H}^3$$

such that σ_s corresponds to $SGr'_s(h, h^*)$ and τ_t corresponds to $SGr'_{s_0}(L_{e^{-it}}(h, h^*))$.

The first-order variations of σ_s and τ_t are the fields

$$(6) \quad X = \frac{\partial \sigma_s}{\partial s}|_{s=s_0}, \quad Y = \frac{\partial \tau_t}{\partial t}|_{t=0}$$

regarded as sections of the fiber-bundle $\Theta = \sigma_{s_0}^*(T\mathbb{H}^3)$ on \tilde{S} . Imposing the equivariance of σ_s under ρ_{is} , and of τ_t under ρ_{t+is_0} , we deduce that

$$(7) \quad X(\tilde{p}) = (\rho(\gamma))_* X(\gamma^{-1}\tilde{p}) + \eta(\gamma)(\tilde{p})$$

$$(8) \quad Y(\tilde{p}) = (\rho(\gamma))_* Y(\gamma^{-1}\tilde{p}) + \xi(\gamma)(\tilde{p})$$

where we have put $\rho = \rho_{z_0}$, $\eta = \eta_{z_0}$, $\xi = \xi_{z_0}$ and we are identifying the elements of $\mathfrak{sl}_2(\mathbb{C})$ with the Killing vector fields on \mathbb{H}^3 .

We will find some explicit relation between X and Y that, used in (7) and (8), shall show equation (5).

5.2. General formulas. We consider any smooth family of immersions

$$\sigma_s : \tilde{S} \rightarrow \mathbb{H}^3$$

and let \tilde{I}_s be the first fundamental form on \tilde{S} and \tilde{B}_s the shape operator associated with σ_s .

Fix $s_0 > 0$: we study σ_s around $s = s_0$. Let us denote by Θ the vector bundle $\sigma_{s_0}^*(T\mathbb{H}^3)$ and let $X = \frac{\partial \sigma}{\partial s}|_{s=s_0}$, seen as a section of Θ . In this section, we will express the derivatives of \tilde{I}_s and \tilde{B}_s at $s = s_0$ in terms of the field X , and we will show that these quantities determine X up to global Killing vector fields.

We notice that there is a natural inclusion of $T\tilde{S}$ into Θ given by $d\sigma_{s_0}$. For the sake of simplicity, we will identify $T\tilde{S}$ with its image in Θ . Given a point $\tilde{p} \in \tilde{S}$ and $s > 0$ we denote by $\tilde{\nu}_s(\tilde{p})$ the unit vector at $\sigma_s(\tilde{p})$ orthogonal to $d\sigma_s(\tilde{p})(T_{\tilde{p}}\tilde{S})$ such that, for every positive basis $\{e_1, e_2\}$ of $T_{\tilde{p}}\tilde{S}$, the vectors $\tilde{\nu}_s(\tilde{p}), d\sigma_s(\tilde{p})(e_1), d\sigma_s(\tilde{p})(e_2)$ form a positively oriented basis of $T_{\sigma_s(\tilde{p})}\mathbb{H}^3$. In this way, if \tilde{J} denotes the complex structure on \tilde{S} (and by abuse of notation on $\sigma_s(\tilde{S})$), we have

$$(9) \quad \tilde{J}_s \tilde{v} = \tilde{\nu}_s \times \tilde{v}$$

for all $\tilde{v} \in d\sigma_s(T\tilde{S})$.

A linear connection D is defined on Θ by pulling back the Levi-Civita connection on $T\mathbb{H}^3$.

The covariant derivative of a section Y of Θ is a linear operator

$$(10) \quad DY : T\tilde{S} \rightarrow T\mathbb{H}^3.$$

Such an operator can be decomposed as the sum of a self-adjoint operator of $T\tilde{S}$ (identified to a subspace of $T\mathbb{H}^3$) and the restriction of a skew-symmetric operator of $T\mathbb{H}^3$.

Lemma 5.3. *Given a section V of Θ , there exist*

- a self-adjoint operator A^V of $T\tilde{S}$;
- a section S^V of Θ

such that

$$(11) \quad D_{\tilde{v}}V = A^V(\tilde{v}) + S^V \times \tilde{v}$$

for every $\tilde{v} \in T\tilde{S}$. Moreover both A^V and S^V are uniquely determined.

Proof. At every point, DV can be decomposed in a tangential part and normal part:

$$D_{\tilde{v}}V = \alpha(\tilde{v}) + \langle D_{\tilde{v}}V, \tilde{v} \rangle \tilde{v}.$$

Clearly α is an operator of $T\tilde{S}$, so it can be decomposed into a self-adjoint part $A(\alpha)$ and a skew-symmetric part $S(\alpha)$. Notice that the skew-symmetric part is a multiple of \tilde{J} , in particular there is $a \in \mathbb{R}$ such that $S(\alpha)(\tilde{v}) = a\tilde{J}(\tilde{v}) = a\tilde{v} \times \tilde{v}$. On the other hand, there exists a tangent vector \tilde{w} such that

$$\langle D_{\tilde{v}}V, \tilde{v} \rangle = \langle \tilde{v}, \tilde{w} \rangle$$

for every $\tilde{v} \in T\tilde{S}$. In particular the normal part of $D_{\tilde{v}}V$ can be regarded as the restriction on $T\tilde{S}$ of the skew-symmetric operator

$$\tilde{v} \mapsto \langle \tilde{v}, \tilde{w} \rangle \tilde{v} - \langle \tilde{v}, \tilde{v} \rangle \tilde{w} = (\tilde{w} \times \tilde{v}) \times \tilde{v}.$$

So if we put $A^V = A(\alpha)$ and $S^V = a\tilde{v} + (\tilde{w} \times \tilde{v})$, Equation (11) is verified.

We show now that this decomposition is unique. Suppose A is a self-adjoint operator of $T\tilde{S}$ and let S be a vector tangent to \mathbb{H}^3 such that $A^V(\tilde{v}) + S^V \times \tilde{v} = A(\tilde{v}) + S \times \tilde{v}$ for all $\tilde{v} \in T\tilde{S}$. Clearly $(A^V - A)(\tilde{v}) = (S - S^V) \times \tilde{v}$. This implies that $S - S^V$ is a normal vector. Since $A - A^V$ is self-adjoint, it follows that $A^V - A = 0$ and $S^V - S = 0$. \square

An important property of the covariant derivative of X is that $D_{\tilde{v}}X$ is the variation of the image of \tilde{v} in \mathbb{H}^3 along the family (σ_s) . More precisely the following statement holds. Here we denote by D/ds the covariant derivative along (σ_s) associated to D .

Lemma 5.4. *Given a tangent vector $\tilde{v} \in T_{\tilde{p}}\tilde{S}$, we consider the field $\tilde{v}_s = d\sigma_s(\tilde{v})$ along the curve $s \mapsto \sigma_s(\tilde{p})$. We have*

$$(12) \quad \frac{D\tilde{v}_s}{ds}|_{s=s_0} = D_{\tilde{v}}X.$$

Proof. Take a path $v : (-\delta, \delta) \rightarrow \tilde{S}$ such that $v'(0) = \tilde{v}$ and consider the map $\tilde{x}(\varepsilon, s) = \sigma_s(v(\varepsilon))$. We have $\tilde{v}_s = \frac{\partial \tilde{x}}{\partial \varepsilon}(0, s)$ whereas $\frac{\partial \tilde{x}}{\partial s}(\varepsilon, s_0) = X(v(\varepsilon))$. A direct computations shows that

$$\frac{D}{ds}\tilde{v}_s|_{s=s_0} = \frac{D}{ds}\frac{\partial \tilde{x}}{\partial \varepsilon}(0, s_0) = \frac{D}{d\varepsilon}\frac{\partial \tilde{x}}{\partial s}(0, s_0) = \frac{D}{d\varepsilon}X(0, s_0) = D_{\tilde{v}}X.$$

\square

Now we apply the decomposition (11) to the field X , so we call A^X the self-adjoint part of DX and X' the field S^X . It turns out that the first order variation of \tilde{I}_s is determined by A^X . On the other hand, the field X' determines the variation of the normal field along the family of σ_s .

Lemma 5.5. *Given $\tilde{u}, \tilde{v} \in T_{\tilde{p}}\tilde{S}$, we have*

$$(13) \quad \frac{d}{ds}\tilde{I}_s(\tilde{u}, \tilde{v})|_{s=s_0} = 2\tilde{I}_s(A^X(\tilde{u}), \tilde{v}), \quad \frac{D\tilde{v}}{ds}|_{s=s_0} = X' \times \tilde{v}.$$

Proof. We have that

$$\tilde{I}_s(\tilde{u}, \tilde{v}) = \langle d\sigma_s(\tilde{u}), d\sigma_s(\tilde{v}) \rangle.$$

Applying Lemma 5.4 we get

$$\frac{d}{ds}\tilde{I}_s(\tilde{u}, \tilde{v}) = \langle D_{\tilde{u}}X, \tilde{v} \rangle + \langle \tilde{u}, D_{\tilde{v}}X \rangle.$$

Since $D_{\tilde{u}}X = A^X(\tilde{u}) + X' \times \tilde{u}$, where A^X is a self-adjoint operator of $T\tilde{S}$, we get the first formula.

To prove the second formula, first we notice that since $\langle \tilde{v}, \tilde{v} \rangle = 1$, then $\frac{D\tilde{v}}{ds}|_{s=s_0}$ is a tangent field. On the other hand, given a tangent vector \tilde{v} we have $\langle \tilde{v}, d\sigma_s(\tilde{v}) \rangle = 0$. Differentiating this identity we get

$$\left\langle \frac{D\tilde{v}}{ds}, \tilde{v} \right\rangle = -\langle \tilde{v}, D_{\tilde{v}}X \rangle = \langle \tilde{v}, X' \times \tilde{v} \rangle.$$

Since $X' \times \tilde{v}$ is tangent, this proves that $\frac{D\tilde{v}}{ds} = X' \times \tilde{v}$. \square

Using (13) we get the first order variation of \tilde{B}_s at $s = s_0$. In the computation of such a variation, the covariant derivative of X' appears. It is useful to apply the decomposition (11) to DX' . In particular, we put $X'' = S^{X'}$, so that

$$D_{\tilde{v}}X' = A^{X'}(\tilde{v}) + X'' \times \tilde{v}.$$

Lemma 5.6. *Given $\tilde{v} \in T_{\tilde{p}}\tilde{S}$, we have*

$$(14) \quad \frac{d}{ds}(\tilde{B}_s(\tilde{v}))|_{s=s_0} = \tilde{J}A^{X'}(\tilde{v}) - \langle X + X'', \tilde{v} \rangle \tilde{v} - A^X \circ \tilde{B}_{s_0}(\tilde{v}).$$

Proof. Differentiating with respect to s the identity

$$d\sigma_s(\tilde{B}_s(\tilde{v})) = -D_{\tilde{v}}\tilde{v}$$

and evaluating at $s = s_0$, we obtain

$$(15) \quad D_{\tilde{B}_{s_0}(\tilde{v})}X + \dot{\tilde{B}}(\tilde{v}) = -\frac{D}{ds}(D_{\tilde{v}}\tilde{v}).$$

On the other hand, we have that

$$(16) \quad \frac{D}{ds}(D_{\tilde{v}}\tilde{v}) = D_{\tilde{v}}\left(\frac{D\tilde{v}}{ds}\right) + \bar{R}(X, \tilde{v})\tilde{v}$$

where \bar{R} is the Riemann tensor of \mathbb{H}^3 .

By (13) we have that

$$(17) \quad D_{\tilde{v}}\left(\frac{D\tilde{v}}{ds}\right) = D_{\tilde{v}}(X' \times \tilde{v}) = A^{X'}(\tilde{v}) \times \tilde{v} + (X'' \times \tilde{v}) \times \tilde{v} - X' \times \tilde{B}_{s_0}(\tilde{v}).$$

On the other hand, since \mathbb{H}^3 has constant curvature -1 , its Riemann tensor is simply given by

$$(18) \quad \bar{R}(X, \tilde{v})\tilde{v} = \langle X, \tilde{v} \rangle \tilde{v} - \langle \tilde{v}, \tilde{v} \rangle X = \langle X, \tilde{v} \rangle \tilde{v}.$$

Using (16), (17), and (18) in (15) we get

$$A^X(\tilde{B}_{s_0}(\tilde{v})) + X' \times \tilde{B}_{s_0}(\tilde{v}) + \dot{\tilde{B}}(\tilde{v}) = -A^{X'}(\tilde{v}) \times \tilde{v} - (X'' \times \tilde{v}) \times \tilde{v} + X' \times \tilde{B}_{s_0}(\tilde{v}) - \langle X, \tilde{v} \rangle \tilde{v}$$

Since $\tilde{v} \times A^{X'}(\tilde{v}) = \tilde{J}A^{X'}(\tilde{v})$ and $(X'' \times \tilde{v}) \times \tilde{v} = \langle X'', \tilde{v} \rangle \tilde{v}$, Equation (14) follows. \square

Finally we show that Equations (13) and (14) determine X up to some global vector field. This is an easy consequence of the following lemma.

Lemma 5.7. *Let V be a section of Θ and let us put $V' = S^V$ and $V'' = S^{V'}$. Suppose that*

$$A^V = 0 \quad \tilde{J}A^{V'} - \langle V'' + V, \tilde{v} \rangle E - A^V \circ \tilde{B}_{s_0} = 0.$$

Then V is the restriction of a global Killing field of \mathbb{H}^3 on \tilde{S} .

Proof. Under the hypothesis of the lemma, neither the induced metric nor the shape operator of the surface vary under the first-order deformation defined by V . The conclusion therefore follows from the Fundamental Theorem of surface theory, see e.g. [31]. \square

5.3. The variation field of SGr' . In this section we apply formulas obtained in the previous subsection to the family of convex immersions $\sigma_s : \tilde{S} \rightarrow \mathbb{H}^3$ defined in (4).

Lemma 5.8. *For $X = \frac{\partial \sigma}{\partial s}|_{s=s_0}$, denote by X' the section S^X and by X'' the section $S^{X'}$. The following formulas hold:*

$$(19) \quad 2A^X = \tanh(s_0/2)E$$

$$(20) \quad A^{X'} = [\tilde{J}, \tilde{b}]/4$$

$$(21) \quad \langle X + X'', \tilde{v} \rangle = \frac{\text{tr}(\tilde{b})}{4}.$$

Proof. The embedding data corresponding to σ_s are

$$\tilde{I}_s = \cosh^2(s/2)\tilde{h} \quad \tilde{B}_s = -\tanh(s/2)\tilde{b}$$

so we easily get that

$$\frac{d}{ds}\tilde{I}_s(\tilde{u}, \tilde{v})|_{s=s_0} = \tanh(s_0/2)\tilde{I}_{s_0}(\tilde{u}, \tilde{v}).$$

Comparing this formula with (13), we get that $2A^X = \tanh(s_0/2)E$.

On the other hand, applying (14) we get

$$-\frac{1}{2 \cosh^2(s_0/2)} \tilde{b} = \tilde{J} A^{X'} - \langle X'' + X, \tilde{\nu} \rangle E + \frac{\tanh^2(s_0/2)}{2} \tilde{b},$$

which can be also written

$$\tilde{b} = -2\tilde{J} A^{X'} + 2\langle X'' + X, \tilde{\nu} \rangle E.$$

Multiplying by \tilde{J} we deduce that

$$\tilde{J}\tilde{b} = 2A^{X'} + 2\langle X'' + X, \tilde{\nu} \rangle \tilde{J}.$$

Notice that this must coincide with the decomposition of $\tilde{J}\tilde{b}$ in symmetric and skew-symmetric part. Since the adjoint of $\tilde{J}\tilde{b}$ is $-\tilde{b}\tilde{J}$ it follows that

$$\begin{aligned} 2A^{X'} &= \frac{\tilde{J}\tilde{b} - \tilde{b}\tilde{J}}{2} = \frac{[\tilde{J}, \tilde{b}]}{2} \\ 2\langle X'' + X, \tilde{\nu} \rangle \tilde{J} &= \frac{\tilde{J}\tilde{b} + \tilde{b}\tilde{J}}{2} = \frac{-\tilde{J}\tilde{b}\tilde{J} + \tilde{b}}{2} \tilde{J} = \frac{\tilde{b}^{-1} + \tilde{b}}{2} \tilde{J} = \frac{\text{tr}(\tilde{b})}{2} \tilde{J} \end{aligned}$$

where we have used that $\tilde{b} + \tilde{b}^{-1} = \text{tr}(\tilde{b})E$. \square

A consequence of Lemma 5.8 is that $X'' + X$ can be explicitly computed.

Proposition 5.9. *With the notation of Lemma 5.8 the following identity holds:*

$$X'' = -X + \frac{\text{tr}(\tilde{b})}{4} \tilde{\nu}.$$

Proof. By (21), it is sufficient to prove that

$$\langle X'' + X, \tilde{\nu} \rangle = 0$$

for every tangent vector $\tilde{\nu}$.

Let \tilde{u}, \tilde{v} be two tangent fields on \tilde{S} . By using the identity

$$D_{\tilde{u}}X = \frac{1}{2} \tanh(s_0/2) \tilde{u} + X' \times \tilde{u}$$

an explicit computation shows that

$$\bar{R}(\tilde{u}, \tilde{v})X = D_{\tilde{u}}D_{\tilde{v}}(X) - D_{\tilde{v}}D_{\tilde{u}}(X) - D_{[\tilde{u}, \tilde{v}]}X = D_{\tilde{u}}X' \times \tilde{v} - D_{\tilde{v}}X' \times \tilde{u}.$$

Moreover,

$$(22) \quad D_{\tilde{u}}X' \times \tilde{v} - D_{\tilde{v}}X' \times \tilde{u} = A^{X'}(\tilde{u}) \times \tilde{v} - A^{X'}(\tilde{v}) \times \tilde{u} + (X'' \times \tilde{u}) \times \tilde{v} - (X'' \times \tilde{v}) \times \tilde{u}.$$

By (20), $A^{X'}$ is a self-adjoint traceless operator, and it follows that the sum of the first two terms of (22) vanishes. Eventually, we get

$$\bar{R}(\tilde{u}, \tilde{v})X = (X'' \times \tilde{u}) \times \tilde{v} - (X'' \times \tilde{v}) \times \tilde{u} = \langle X'', \tilde{v} \rangle \tilde{u} - \langle X'', \tilde{u} \rangle \tilde{v}.$$

Since $\bar{R}(\tilde{u}, \tilde{v})X = \langle X, \tilde{u} \rangle \tilde{v} - \langle X, \tilde{v} \rangle \tilde{u}$, we easily deduce that $\langle X'' + X, \tilde{\nu} \rangle = 0$ for all tangent vectors $\tilde{\nu}$. \square

Proposition 5.10. *Let $\tau_t : \tilde{S} \rightarrow \mathbb{H}^3$ be the family of convex immersions corresponding to the projective surface $SGr'_{s_0} \circ L_{e^{-it}}(h, h^*)$ and denote by Y its first order variation at $t = 0$. Then $Y = -X'$ up to adding a global Killing vector field, where X' is the vector field defined in Lemma 5.8.*

Proof. Let \tilde{I}_t be the first fundamental form corresponding to τ_t and let \tilde{B}_t be the corresponding shape operator. According to Lemmas 5.5, 5.6 and 5.7, it is sufficient to show that

$$(23) \quad \frac{d}{dt} \tilde{I}_t(\tilde{u}, \tilde{v})|_{t=0} = -2\tilde{I}_0(A^{X'}(\tilde{u}), \tilde{v})$$

$$(24) \quad \frac{d}{dt} \tilde{B}_t(\tilde{u})|_{t=0} = -\left(\tilde{J}A^{X''}(\tilde{u}) - \langle X' + X''', \tilde{\nu} \rangle \tilde{u} - A^{X'} \circ \tilde{B}_0(\tilde{u})\right).$$

where $X''' = S^{X''}$ is the vector field corresponding to the skew-symmetric part of DX'' . Call $\beta_t = \cos(t/2)E - \sin(t/2)Jb$, so that we have

$$(25) \quad I_t = \cosh^2(s_0/2)h(\beta_t \bullet, \beta_t \bullet)$$

$$(26) \quad B_t = -\tanh(s_0/2)\beta_{-t}b\beta_t.$$

It follows that

$$\begin{aligned} \frac{d}{dt} \tilde{I}_t(\tilde{u}, \tilde{v})|_{t=0} &= \frac{1}{2} \left(-\tilde{I}_0(\tilde{J}\tilde{b}(\tilde{u}), \tilde{v}) - \tilde{I}_0(\tilde{u}, \tilde{J}\tilde{b}(\tilde{v})) \right) \\ &= -\frac{1}{2} \tilde{I}_0([\tilde{J}, \tilde{b}]\tilde{u}, \tilde{v}) = -2\tilde{I}_0(A^{X'}(\tilde{u}), (\tilde{v})). \end{aligned}$$

To show equation (24), we first compute DX'' . By Proposition 5.9 we have

$$\begin{aligned} D_u X'' &= -A^X u - X' \times u - \frac{\text{tr}(\tilde{b})}{4} \tilde{B}_0 u + \left\langle \text{grad} \left(\frac{\text{tr}(\tilde{b})}{4} \right), u \right\rangle \tilde{v} \\ &= - \left(A^X + \frac{\text{tr}(\tilde{b})}{4} \tilde{B}_0 \right) u - (X' + \tilde{J} \text{grad}(\text{tr}(\tilde{b})/4)) \times u. \end{aligned}$$

where grad is the gradient on \tilde{S} with respect to \tilde{I}_0 . In particular,

$$X''' = -X' - \tilde{J} \text{grad}(\text{tr}(\tilde{b})/4), \quad 2A^{X''} = -\tanh(s_0/2) \left(E - \frac{\text{tr}(\tilde{b})}{2} \tilde{b} \right).$$

Replacing these identities in the right hand side of (24), we deduce

$$\begin{aligned} 2(\tilde{J}A^{X''}(\tilde{u}) - \langle X' + X''', \tilde{v} \rangle \tilde{u} - A^{X'} \circ \tilde{B}_0(\tilde{u})) &= -\tanh(s_0/2) \left(\tilde{J} - \frac{\text{tr}(\tilde{b})}{2} \tilde{J}\tilde{b} \right)(\tilde{u}) + \tanh(s_0/2) \frac{(\tilde{J}\tilde{b} - \tilde{b}\tilde{J})}{2} \tilde{b}(\tilde{u}) \\ &= -\tanh(s_0/2) \left(\tilde{J} - \frac{\text{tr}(\tilde{b})}{2} \tilde{J}\tilde{b} - \frac{1}{2} \tilde{J}\tilde{b}^2 + \frac{\tilde{J}}{2} \right)(\tilde{u}) \end{aligned}$$

Using the identity $\tilde{b}^2 = \text{tr}(\tilde{b})\tilde{b} - E$, we obtain that the right hand side in (24) is equal to

$$(27) \quad \tanh(s_0/2) (\tilde{J} - (\text{tr}(\tilde{b})/2) \tilde{J}\tilde{b})$$

On the other hand, (26) shows that the left hand side of (24) is equal to

$$(28) \quad \frac{\tanh(s_0/2)}{2} (\tilde{b}\tilde{J}\tilde{b} - \tilde{J}\tilde{b}^2) = \frac{\tanh(s_0/2)}{2} (\tilde{J} - \tilde{J}(\text{tr}(\tilde{b})\tilde{b} - E)) = \frac{\tanh(s_0/2)}{2} (2\tilde{J} - \text{tr}(\tilde{b})\tilde{J}\tilde{b}).$$

Equation (24) follows by comparing (27) with (28). \square

5.4. The comparison of the cocycles. Any element of $K \in \mathfrak{sl}_2(\mathbb{C})$ can be regarded as a Killing vector field on \mathbb{H}^3 . Notice that by definition of Killing vector field, there is another field K' associated to K such that

$$D_{\tilde{v}} K = K' \times \tilde{v}.$$

Using the same argument as in Proposition 5.9, one can check that K' is a Killing vector field and in fact $K'' = -K$. More precisely, we have the following lemma:

Lemma 5.11. *As elements of $\mathfrak{sl}_2(\mathbb{C})$ we have $K' = iK$.* \square

Given a point $\tilde{x} \in \mathbb{H}^3$ we have a natural map

$$\text{ev}_{\tilde{x}} : \mathfrak{sl}_2(\mathbb{C}) \ni K \mapsto (K(\tilde{x}), K'(\tilde{x})) \in T_{\tilde{x}}\mathbb{H}^3 \oplus T_{\tilde{x}}\mathbb{H}^3.$$

It is a well-known fact that such a map is an isomorphism for every $\tilde{x} \in \mathbb{H}^3$.

Because of Lemma 5.11, if $\text{ev}_{\tilde{x}}(K) = (\tilde{w}_1, \tilde{w}_2)$, then $\text{ev}_{\tilde{x}}(iK) = (\tilde{w}_2, -\tilde{w}_1)$.

Given any section V on Θ , we define

$$K^V : \tilde{S} \rightarrow \mathfrak{sl}_2(\mathbb{C})$$

such that $\text{ev}_{\tilde{x}}(K^V(\tilde{p})) = V(\tilde{p})$ and $\text{ev}_{\tilde{x}}((K^V)'(\tilde{p})) = V'(\tilde{p})$ for every $\tilde{x} = \sigma_{s_0}(\tilde{p})$.

In particular, we have maps K^X and K^Y associated to the fields X, Y defined in (6), so that $\text{ev}_{\tilde{x}}(K^X) = (X, X')$ and $\text{ev}_{\tilde{x}}(K^Y) = (-X', -X'')$ by Proposition 5.10. We conclude by Proposition 5.9 that

$$K^Y = -iK^X - K_0$$

where $K_0(\tilde{p}) = \text{ev}_{\tilde{x}}^{-1}(0, X'' + X) = \text{ev}_{\tilde{x}}^{-1}(0, \frac{\text{tr}(\tilde{b})}{4} \tilde{v})$ for all $\tilde{x} = \sigma_{s_0}(\tilde{p})$. Since the field $\frac{\text{tr}(\tilde{b})}{4} \tilde{v}$ is invariant under the action of $\pi_1(S)$, we find that K_0 is equivariant, that is

$$K_0(\gamma\tilde{p}) = \text{Ad}(\rho(\gamma))K_0(\tilde{p}).$$

However, it follows from Equation (7) that

$$(29) \quad K^X(\gamma\tilde{p}) - \text{Ad}(\rho(\gamma))K^X(\tilde{p}) = \eta(\gamma)$$

$$(30) \quad K^Y(\gamma\tilde{p}) - \text{Ad}(\rho(\gamma))K^Y(\tilde{p}) = \xi(\gamma)$$

and by these equations and identities (7) and (8) one deduces that

$$\xi(\gamma) = -i\eta(\gamma) - (\text{Ad}(\rho(\gamma)) - 1)K_0(\tilde{p}) = -i\eta(\gamma).$$

Thus (5) is proved.

5.5. Parameterization by the disk. The parameterization of the complex landslide used above is well-suited for a comparison with the complex earthquake. However, another parameterization – already used in the introduction – is perhaps more convenient when considering the holomorphic disks in Teichmüller space obtained as the image of the complex flow. This new parameter ζ takes values in the unit disk. We develop here the relationship between these two parameterizations and we investigate the regularity at $\zeta = 0$.

Consider $z = t + is$ in the upper half-plane, so that $t \in \mathbb{R}$ and $s \geq 0$, and we set $\zeta = \exp(iz) = \exp(-s + it)$, which belongs to the punctured closed unit disk $\bar{\Delta} = \{\zeta \in \mathbb{C} \mid 0 < |\zeta| \leq 1\}$. For $h, h^* \in \mathcal{T}$ we then define

$$P_\zeta(h, h^*) := P'_{t+is}(h, h^*) \quad C_\zeta(h, h^*) := C'_{t+is}(h, h^*) .$$

These maps are well-defined since P' and C' are invariant under $t \mapsto t + 2\pi$. Clearly, the maps $\zeta \mapsto P_\zeta(h, h^*)$ and $\zeta \mapsto C_\zeta(h, h^*)$ are holomorphic in the unit disk minus its center, for any fixed h and h^* .

We first give an explicit formula for $C'_{is}(h, h^*)$.

Lemma 5.12. *Let $h, h^* \in \mathcal{T}$ be two hyperbolic metrics on S , and let b be the bundle morphism appearing in Corollary 1.5. For every $s \in \mathbb{R}_{\geq 0}$,*

$$C'_{is}(h, h^*) = h(\gamma_s \bullet, \gamma_s \bullet)$$

as conformal structures, where

$$\gamma_s = \cosh(s/2)E + \sinh(s/2)b .$$

Proof. By definition, $C'_{is}(h, h^*)$ is the conformal structure at infinity of the (unique) hyperbolic end containing a convex surface with induced metric $I = \cosh^2(s/2)h$ and third fundamental form $\text{III} = \sinh^2(s/2)h^*$. Its shape operator is then $B = \tanh(s/2)b$, and conformal structure at infinity of the end is given (see e.g. [28]) by

$$C'_{is}(h, h^*) = I((E + B) \bullet, (E + B) \bullet) = h(\gamma_s \bullet, \gamma_s \bullet) .$$

□

We can now give a general formula for $C'_{t+is}(h, h^*)$ for $t + is \in \bar{\mathbb{H}}$.

Lemma 5.13. *Let $s, t \in \mathbb{R}$ with $s \geq 0$, and let $\zeta = \exp(-s + it)$. Then, for all $h, h^* \in \mathcal{T}$,*

$$C_\zeta(h, h^*) = h(B_\zeta^\# \bullet, B_\zeta^\# \bullet) ,$$

where

$$B_\zeta^\# = \frac{\zeta + 1}{2\sqrt{\zeta}}E - \frac{\zeta - 1}{2\sqrt{\zeta}}b$$

and $\sqrt{\zeta}$ is a notation for $\exp((-s + it)/2)$.

Here we use the convention that the complex number i acts as the complex structure J on tangent vectors.

Proof. It follows from the definition of C' and from the previous lemma that

$$C'_{t+is}(h, h^*) = h(\beta_{-t} \circ \overline{\gamma}_s \bullet, \beta_{-t} \circ \overline{\gamma}_s \bullet) ,$$

where (as in Section 3):

$$\beta_t = \cos(t/2)E + \sin(t/2)Jb , \quad \overline{\gamma}_s = \cosh(s/2)E + \sinh(s/2)b_{-t} , \quad b_{-t} = \beta_t \circ b \circ \beta_{-t} .$$

It is then clear that

$$\beta_{-t} \circ \overline{\gamma}_s = \gamma_s \circ \beta_{-t} ,$$

so that

$$C'_{t+is}(h, h^*) = h(B_\zeta^\# \bullet, B_\zeta^\# \bullet) ,$$

where $B_\zeta^\# = \gamma_s \circ \beta_{-t}$.

Using the fact that $bJbJ = -E$:

$$\begin{aligned}
B_\zeta^\# &= (\cosh(s/2)E + \sinh(s/2)b) \circ (\cos(t/2)E - \sin(t/2)Jb) \\
&= (\cosh(s/2)\cos(t/2)E - \sinh(s/2)\sin(t/2)bJb) + (\cos(t/2)\sinh(s/2)E - \cosh(s/2)\sin(t/2)J)b \\
&= (\cosh(s/2)\cos(t/2)E - \sinh(s/2)\sin(t/2)J) + (\cos(t/2)\sinh(s/2)E - \cosh(s/2)\sin(t/2)J)b \\
&= \cosh((-s+it)/2)E - \sinh((-s+it)/2)b.
\end{aligned}$$

Setting $\sqrt{\zeta} = \exp((-s+it)/2)$, we can write this relation as

$$B_\zeta^\# = \frac{\sqrt{\zeta} + 1/\sqrt{\zeta}}{2}E - \frac{\sqrt{\zeta} - 1/\sqrt{\zeta}}{2}b.$$

□

It follows from the definitions that C is essentially the same as C' with a different parameterization. The main properties of this map are as follows.

Proposition 5.14. *Let $h, h^* \in \mathcal{T}$ and let c be the “center” of (h, h^*) as defined in Section 1.6. Then:*

- (1) $C_\zeta(h, h^*)$ is defined for all $\zeta \in \bar{\Delta}$,
- (2) it is holomorphic in ζ ,
- (3) it extends continuously, and therefore holomorphically, at $\zeta = 0$, with $C_0(h, h^*) = c$.
- (4) it also extends holomorphically to the open disk of center 0 and radius $(\kappa_0 + 1)/(\kappa_0 - 1)$, where $\kappa_0 = \max_{x \in S} \kappa(x)$ and $\kappa : S \rightarrow [1, \infty)$ is the bigger eigenvalue of the operator b associated to the minimal Lagrangian map isotopic to the identity between (S, h) and (S, h^*) (see Section 1.3).

In particular, c appears as a smooth point of the holomorphic disk defined by C , while it was obtained only in the limit $s \rightarrow \infty$ in the parameterization used by C' .

Remark 5.15. Unlike the map $C_\bullet(h, h^*)$, the map $P_\bullet(h, h^*)$ does not extend at $\zeta = 0$. Indeed, take any sequence of positive real numbers $\zeta_n \rightarrow 0$. By definition of the map P , there is an embedding of S into the hyperbolic end M_n , with first fundamental form equal to $I_n = \cosh^2(-\frac{1}{2} \log \zeta_n)h$ and shape operator $B_n = -\tanh(-\frac{1}{2} \log \zeta_n)b$, which corresponds to the projective structure P_n . In particular, B_n converges to $-b$. On the other hand, by Proposition 4.2 of [15], if P_n converges to a projective surface, B_n should converge to the identity.

Proof of Proposition 5.14. The first two points are direct consequences of the definition of C from C' , and of Theorem 5.1. The third point follows from the expression of $B_\zeta^\#$ in Lemma 5.13, because $C_\zeta(h, h^*)$ is really considered as a conformal structure, so it is not changed if we multiply $B_\zeta^\#$ by a complex-valued function defined on S . In particular, we can multiply $B_\zeta^\#$ by $2\sqrt{\zeta}$, obtaining

$$2\sqrt{\zeta}B_\zeta^\# = (1 + \zeta)E + (1 - \zeta)b,$$

which is clearly continuous at $u = 0$.

For the last point note that the expression defining $C_\zeta(h, h^*)$ in Lemma 5.13 can be analytically continued if $B_\zeta^\#$ is non-singular at all points of S . This happens if

$$\frac{\zeta + 1}{2\sqrt{\zeta}} + \frac{1 - \zeta}{2\sqrt{\zeta}}\kappa \neq 0$$

everywhere on S , which is certainly satisfied if

$$|\zeta| < \frac{\kappa_0 + 1}{\kappa_0 - 1}.$$

□

6. THE EARTHQUAKE FLOW AS A LIMIT

The main goal of this section is to prove Theorem 1.12. The arguments are based on comparing surfaces in AdS_3 with constant Gauss curvature close to -1 to pleated surfaces. The key step in the proof of Theorem 1.12 will be Theorem 1.13.

We fix a hyperbolic metric h on S and a divergent sequence of metrics $h_n^* \in \mathcal{T}$. We will study the asymptotic behavior of the holomorphic map $z \mapsto P_z(h, h_n^*)$ assuming that $(h_n^*)_{n \in \mathbb{N}}$ converges to a point in the Thurston boundary of $\mathcal{T}(S)$ which is the projective class of some measured geodesic lamination λ on S .

Take any sequence $\theta_n > 0$ with $\lim_{n \rightarrow \infty} \theta_n = 0$ such that $\theta_n \ell_{h_n^*} \rightarrow \iota(\lambda, \bullet)$, i.e. for every free homotopy class γ of closed curves of S , the h_n^* -length of the h_n^* -geodesic representative of γ rescaled by the factor θ_n converges

to the intersection between γ and λ . Define $P'_n : \overline{\mathbb{H}} \rightarrow \mathcal{P}(S)$ as $P'_n(z) = P'_{\theta_n z}(h, h_n^*)$, which are holomorphic, and let $P'_\infty : \overline{\mathbb{H}} \rightarrow \mathcal{P}(S)$ be $P'_\infty(t + is) = Gr_{s\lambda/2}(E_{-t\lambda/2}(h))$.

Theorem 6.1. *For every $z \in \overline{\mathbb{H}}$, we have that $P'_n(z) \rightarrow P'_\infty(z)$.*

Notice that since the P'_n are holomorphic, the convergence $P'_n \rightarrow P'_\infty$ is, in fact, in C^∞ . Note also that in this section we use the parameterization by the upper half-plane – which is more practical when considering the limit to complex earthquakes – so that we use the notations with primes for L, P, SGr , etc.

6.1. Convergence on the imaginary axis. In this subsection we prove that if $\theta_n \ell_{h_n^*} \rightarrow \iota(\lambda, \bullet)$, then

$$(31) \quad P'_{i\theta_n}(h, h_n^*) \rightarrow Gr_{\lambda/2}(h).$$

In fact, in order to prove Theorem 6.1, we will need to prove that the convergence is uniform with respect to h .

Proposition 6.2. *Let $(h_n)_{n \in \mathbb{N}}$ be a sequence of hyperbolic metrics converging to a hyperbolic metric h on S , and let $(h_n^*)_{n \in \mathbb{N}}$ be a sequence of hyperbolic metrics converging projectively to $[\lambda]$ in the Thurston boundary of \mathcal{T} . If θ_n is a sequence of positive numbers such that $\theta_n \ell_{h_n^*} \rightarrow \iota(\lambda, \bullet)$, then $SGr'_{\theta_n}(h_n, h_n^*)$ converges to $Gr_{\lambda/2}(h)$.*

Notice that (31) corresponds to the particular case of Proposition 6.2 in which h_n is constant.

By definition, $SGr'_{\theta_n}(h_n, h_n^*) = P'_{i\theta_n}(h_n, h_n^*)$ is the projective structure on S determined by prescribing that the associated hyperbolic end (M_n, g_{M_n}) contains a constant curvature surface S_n , parametrized by $\bar{\sigma}_n : S \rightarrow S_n \subset M_n$, with first fundamental form $I_{S_n} = \cosh^2(\theta_n/2)h_n$ and third fundamental form $\mathbb{I}_{S_n} = \sinh^2(\theta_n/2)h_n^*$.

In general, given an embedding $S \rightarrow \Sigma \subset X$ of S inside a (hyperbolic, de Sitter or anti de Sitter) 3-manifold, we will denote by the same symbol the first fundamental form I_Σ (resp. the third fundamental form \mathbb{I}_Σ) and its pull-back on S .

Let ∂M_n be the hyperbolic boundary of M_n , that carries a hyperbolic induced metric $g_{\partial M_n}$ and is locally bent along a measured geodesic lamination λ_n . By definition, $P'_{i\theta_n}(h_n, h_n^*) = Gr_{\lambda_n}(g_{\partial M_n})$. So, in order to prove Proposition 6.2, it is sufficient to check that $(S, g_{\partial M_n})$ converges to (S, h) , and λ_n converges to $\lambda/2$ in $\mathcal{ML}(S)$.

Lemma 6.3. *The hyperbolic metrics $g_{\partial M_n}$ are contained in a compact subset of \mathcal{T} .*

Proof. The closest point projection $r_n : M_n \rightarrow \partial M_n$ is 1-Lipschitz. In particular, we have a 1-Lipschitz map $r_n|_{S_n} : S_n \rightarrow \partial M_n$. This implies that the marked length spectrum of ∂M_n is bounded from above by the marked length spectrum of S_n , that in turn is locally uniformly bounded. \square

Lemma 6.4. *The bending laminations λ_n are contained in a compact subset of \mathcal{ML} .*

In order to prove Lemma 6.4, we will consider the de Sitter spacetime M_n^* dual to M_n : it is the set of complete geodesic planes contained in M_n . The de Sitter structure is induced by the natural map $dev^* : \tilde{M}_n^* \rightarrow d\mathbb{S}^3$, where the model of de Sitter geometry is the set of geodesic planes of \mathbb{H}^3 (see [25, 5]).

Scannell [25] showed that M_n^* is a globally hyperbolic spacetime diffeomorphic to $S \times \mathbb{R}$. Following [5], \tilde{M}_n^* has a natural boundary corresponding to the set of support planes of $\partial \tilde{M}_n$. This boundary is called the initial singularity: the de Sitter metric extends to the boundary and makes it an achronal (but not spacelike) surface. So $\partial \tilde{M}_n^*$ carries a pseudometric d_0 induced by \tilde{M}_n^* . By [5, 6] it turns out that the action of $\pi_1(S)$ on \tilde{M}_n^* extends to the boundary (even if the action on the boundary is neither proper nor free) and the marked length spectrum of this action coincides with the intersection with the bending lamination:

$$(32) \quad \iota(\lambda_n, \gamma) = \inf\{d_0(x^*, \gamma x^*) | x^* \in \partial \tilde{M}_n^*\}.$$

Let S_n^* be the surface in M_n^* dual to S_n , corresponding to the set of support planes of S_n (that by the convexity of S_n are complete planes in M_n). There is a natural map $S_n \rightarrow S_n^*$ sending x to the dual of the plane tangent to S_n at x .

A simple local computation shows that the first fundamental form of S_n^* coincides with the third fundamental form of S_n (through the natural map $S_n \rightarrow S_n^*$). In particular S_n^* is a surface of constant curvature $-1/\sinh^2(\theta_n/2)$. Barbot, Beguin and Zeghib [3] have shown that there is a time function (K-time) $t_n : M_n^* \rightarrow (-\infty, 0)$ such that $t_n^{-1}(k)$ is the unique surface in M_n^* with constant curvature k .

Lemma 6.4 is a simple consequence of (32) and the following general Lemma of de Sitter geometry.

Lemma 6.5 ([4]). *Let M^* be a de Sitter MGHC spacetime and $\partial \tilde{M}^*$ be the boundary of its universal covering. If S^* is a constant curvature surface in M^* , there is a natural 1-Lipschitz equivariant map*

$$\tilde{r}^* : \tilde{S}^* \rightarrow \partial \tilde{M}^*$$

such that $\tilde{r}^(\tilde{x}) \in I^-(\tilde{x}) \cap \partial \tilde{M}^*$.*

In particular, we have

$$\ell_{\partial M^\star}(\gamma) \leq \ell_{S^\star}(\gamma)$$

where $\ell_{\partial M^\star}$ and ℓ_{S^\star} are the marked length spectra of ∂M^\star and S^\star respectively.

Proposition 6.6. *The hyperbolic metrics $g_{\partial M_n}$ converge to h .*

Proof. By Lemmas 6.3 and 6.4, we have that up to passing to a subsequence $M_n \rightarrow M_\infty$. In particular, we can concretely realize $M_n \cong (S \times [0, +\infty), g_{M_n})$ in such a way that g_{M_n} converges to a hyperbolic metric g_{M_∞} such that $(S \times [0, +\infty), g_{M_\infty}) \cong M_\infty$, where we have denoted by the same symbol the metric g_{M_n} on M_n and the corresponding metric on the model $S \times [0, +\infty)$.

By abuse of notation, we denote again by $r_n : S \times [0, +\infty) \rightarrow S \times \{0\}$ the retraction corresponding to the retraction of M_n onto ∂M_n . Let $\bar{\sigma}_n : (S, h) \rightarrow (S \times [0, +\infty), g_{M_n})$ be the embedding with first fundamental form $I_{S_n} = \cosh^2(\theta_n/2)h_n$ and third fundamental form $\mathbb{I}_{S_n} = \sinh^2(\theta_n/2)h_n^\star$.

Notice that the composition $i_n = r_n \circ \bar{\sigma}_n : (S, \cosh^2(\theta_n/2)h_n) \rightarrow (S \times \{0\}, g_{\partial M_n})$ is a 1-Lipschitz homotopy equivalence. So i_n converges (up to passing to a subsequence) to a 1-Lipschitz homotopy equivalence $i_\infty : (S, h) \rightarrow (S \times \{0\}, g_{\partial M_\infty})$. Since both h and $g_{\partial M_\infty}$ are hyperbolic metrics, we conclude that i_∞ is an isometry. \square

Let λ_∞ the bending lamination of M_∞ . In order to conclude the proof of (31) we need to show that $\lambda_\infty = \lambda/2$. In fact, the following general result in Lorentzian geometry and (32) show that

$$(33) \quad \iota(\lambda_\infty, \gamma) = \lim_n \ell_{S_n^\star}(\gamma) = \lim_n \ell_{\mathbb{I}_{S_n}}(\gamma) = \lim_n \frac{\theta_n}{2} \ell_{h_n^\star}(\gamma) = \iota(\lambda/2, \gamma).$$

for every closed curve γ .

Proposition 6.7. [4] *Let $(X_n^\star)_{n \in \mathbb{N}}$ be a sequence of MGH de Sitter (or anti de Sitter) spacetimes homeomorphic to $S \times \mathbb{R}$. Suppose that X_n^\star converges to a MGH spacetime X_∞^\star . Take any sequence of numbers $k_n \rightarrow -\infty$ and let Σ_n^\star be the future-convex surface of constant curvature k_n contained in X_n^\star .*

Denote by ℓ_0 the length spectrum of the initial singularity of X_∞^\star . Then, for every $\gamma \in \pi_1(S)$ we have

$$\ell_{\Sigma_n^\star}(\gamma) \rightarrow \ell_0(\gamma)$$

as $n \rightarrow +\infty$.

In the next section we will give a short description of the initial singularity for anti de Sitter spacetimes and we will apply Proposition 6.7 to this case.

6.2. Convergence on the real axis. In this section we will prove that

$$(34) \quad L_{\theta_n}^1(h, h_n^\star) \rightarrow E_{\lambda/2}(h)$$

Recall that $L_{\theta_n}^1$ is the composition of the map $L'_{\theta_n} : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T} \times \mathcal{T}$ with the projection on the first factor. As in the previous section, we will need a slightly stronger statement.

Proposition 6.8. *Let $(h_n)_{n \in \mathbb{N}}$ and $(h_n^\star)_{n \in \mathbb{N}}$ be two sequences of hyperbolic metrics such that $(h_n)_{n \in \mathbb{N}}$ converges to a hyperbolic metric h on S and that $(h_n^\star)_{n \in \mathbb{N}}$ converges to a point $[\lambda]$ in Thurston boundary of \mathcal{T} . If θ_n is a sequence of positive numbers such that $\theta_n \ell_{h_n^\star} \rightarrow \iota(\lambda, \bullet)$, then $L_{\theta_n}^1(h_n, h_n^\star)$ converges to $E_{\lambda/2}(h)$.*

Recall that the holonomy of $L_{\theta_n}^1(h_n, h_n^\star)$ corresponds to the left holonomy of the MGH AdS $N_n \cong S \times \mathbb{R}$ containing a future-convex K -surface F_n with $I_{F_n} = \cos^2(\theta_n/2)h_n$ and $\mathbb{I}_{F_n} = \sin^2(\theta_n/2)h_n^\star$. In order to prove Proposition 6.8, we will show that N_n converges to a MGH AdS structure N_∞ and F_n converges to the lower boundary $\partial_- C(N_\infty)$ of the convex core of N_∞ . Then we will prove that $\partial_- C(N_\infty)$ is isometric to (S, h) and is bent along a lamination corresponding to $\lambda/2$. By a result of Mess [20], the left holonomy of N_∞ (that is, by definition, the limit of the left holonomies of N_n) is equal to the holonomy of $E_{\lambda/2}(h)$, and Proposition 6.8 follows.

In order to prove that N_n converges to some structure, we will consider the lifting $\phi_n : \tilde{S} \rightarrow \tilde{N}_n \subset \text{AdS}^3$ corresponding to the embedding $\bar{\phi}_n : S \rightarrow F_n \subset N_n$. The map ϕ_n is determined up to isometry of AdS^3 and we will normalize it by requiring that, for some fixed $\tilde{p}_0 \in \tilde{S}$, $\phi_n(\tilde{p}_0) = \tilde{x}_0$ and the normal vector to $\tilde{F}_n = \phi_n(\tilde{S})$ at \tilde{x}_0 is equal to \tilde{v}_0 for some fixed \tilde{x}_0, \tilde{v}_0 in AdS^3 .

The first step to prove the convergence of N_n is to show that ϕ_n converges to a spacelike embedding into AdS^3 .

Proposition 6.9. *Up to passing to a subsequence, \tilde{F}_n converges to a spacelike surface \tilde{F}_∞ in AdS^3 and the map ϕ_n converges to an embedding*

$$\phi_\infty : \tilde{S} \rightarrow \text{AdS}^3$$

whose image is \tilde{F}_∞ .

The easy part of the proof is to show that \tilde{F}_n converges to an embedded surface \tilde{F}_∞ in AdS^3 that is achronal. The main issue is to show that the surface \tilde{F}_∞ is spacelike. The proof relies on the fact that, for some fixed $\tilde{p} \in \tilde{S}$, the tangent planes of \tilde{F}_n at $\phi_n(\tilde{p})$ are uniformly spacelike, in the sense that they cannot approximate lightlike planes.

The proof of this fact is based on the technical Lemma 6.10.

Lemma 6.10. *Let b_n be the h_n -self adjoint operator such that $h_n^* = h_n(b_n \bullet, b_n \bullet)$ and let $\bar{\sigma}_n : S \rightarrow S_n \subset M_n$ be the embedding introduced in Section 6.1. Let $I_{S_n}^\#$ be the lifting to \tilde{S} of the grafted metric $I_{S_n}^\#$ introduced in Definition 2.1. Then, for every compact set $K \subset \tilde{S}$, there is a constant C_K such that the diameter of K with respect to $I_{S_n}^\#$ is bounded by C_K for every n .*

Proof. For any $k \in [-1, 0)$, M_n contains exactly one K -surface of constant curvature k , denoted here by $M_n(k)$ (where by $M_n(-1)$ we mean the boundary of M_n). For each n , let $G_n = SGr'_n(h_n, h_n^*)$ be the projective surface at infinity of M_n . Let us consider the natural retraction $\Pi_{M_n(k)} : G_n \rightarrow M_n(k)$, which is the limit of the closest point projections $M_n(K) \rightarrow M_n(k)$ onto the convex surface $M_n(k)$ as $K > k$ converges to 0 (see Figure 1).




FIGURE 1. The retraction $\Pi_{M_n(k)}$.

On the universal covering, $\Pi_{\tilde{M}_n(k)}$ sends a point $\tilde{x} \in \tilde{G}_n$ to the tangency point of the unique horocycle centered at \tilde{x} and tangent to $\tilde{M}_n(k)$. For $k > -1$, $\Pi_{M_n(k)}$ is a diffeomorphism and the inverse is the map obtained by sending each point of $y \in M_n(k)$ to the final point of the geodesic ray starting from y and orthogonal to $M_n(k)$. If $k = -1$, $\Pi_{M_n(-1)}$ is not injective in general, since points on $M_n(-1)$ can admit several normal directions. Nevertheless, $\Pi_{M_n(-1)} : G_n \rightarrow M_n(-1)$ is a homotopy equivalence.

In [28], it has been showed that this diffeomorphism is conformal with respect to the grafted metric $I_{M_n(k)}^\#$ of $M_n(k)$. The conformal factor is an increasing function of k : this precisely means that the conformal map

$$\Pi_{M_n(k')} \circ \Pi_{M_n(k)}^{-1} : (M_n(k), I_{M_n(k)}^\#) \rightarrow (M_n(k'), I_{M_n(k')}^\#)$$

decreases the lengths when $k > k'$.

Now notice that S_n is equal to $M_n(K_n)$ for $K_n = -1/\cosh^2(\theta_n)$. As definitively $K_n < -1/2$, the map

$$j_n = \bar{\sigma}_n^{-1} \circ \Pi_{S_n} \circ \Pi_{M_n(-1/2)}^{-1} : (M_n(-1/2), I_{M_n(-1/2)}^\#) \rightarrow (S, I_{S_n}^\#)$$

decreases the lengths.

Since M_n converges to an hyperbolic end M_∞ , the surface $M_n(-1/2)$ converges to $M_\infty(-1/2)$ in C^∞ -sense. This means that M_n can be concretely realized as a hyperbolic metric g_{M_n} on $S \times [0, +\infty)$ such that $M_n(-1/2) = S \times \{1\}$ and such that g_{M_n} converges to a hyperbolic metric g_{M_∞} and $M_\infty(-1/2) = S \times \{1\}$.

Then the family of 1-Lipschitz maps j_n converges to the map $j_\infty = \bar{\sigma}_\infty^{-1} \circ \Pi_{S_\infty} \circ \Pi_{M_\infty(-1/2)}^{-1} : M_\infty(-1/2) \rightarrow S$, which is a homotopy equivalence.

Let \tilde{j}_∞ and \tilde{j}_n be the lifting of those maps to the universal covering. Notice that \tilde{j}_∞ is a proper map. If K is a compact subset of \tilde{S} , then $K' = \tilde{j}_\infty^{-1}(K)$ is a compact subset of $\tilde{S} \times \{1\}$, and $K'_n = \tilde{j}_n^{-1}(K)$ is contained in some compact neighborhood of K' for every n . In particular, there exists a constant C'_K such that the diameter of K'_n with respect to $\tilde{I}_{M_n(-1/2)}^\#$ is bounded by C'_K for all n . Taking C_K bigger than C'_K , it follows that the diameter of every K'_n with respect to $\tilde{I}_{M_n(-1/2)}^\#$ is bounded by C_K . Since \tilde{j}_n decreases the lengths, we have that the diameter of K with respect to $I_{\tilde{S}_n}^\#$ is bounded by C_K for n large enough. \square

Lemma 6.11. *For every $d > 0$ there is a compact set K in AdS^3 such that for $\tilde{p} \in \tilde{S}$ with $d_h(\tilde{p}, \tilde{p}_0) < d$, the normal vector $\tilde{\nu}_n(\tilde{p})$ of \tilde{F}_n at $\phi_n(\tilde{p})$ lies in K .*

Proof. Lemma 6.10 implies that for any $d > 0$, there is $D > 0$ such that for any n and any $\tilde{p} \in B_{\tilde{h}}(\tilde{p}_0, d)$ there exists a path $\tilde{\zeta} : [0, 1] \rightarrow \tilde{S}$ connecting \tilde{p}_0 to \tilde{p} such that $\ell_{I_{\tilde{S}_n}^\#}(\tilde{\zeta})$ is bounded by D .

We claim (and will prove below) that this implies that

$$(35) \quad |\langle \tilde{x}_0, \tilde{\nu}_n(\tilde{p}) \rangle| \leq 2e^{2D}$$

$$(36) \quad |\langle \tilde{\nu}_0, \tilde{\nu}_n(\tilde{p}) \rangle| \leq 2e^{2D}.$$

It follows from this claim that $\tilde{\nu}_n(\tilde{p})$ is contained in

$$K = \{w \in \text{AdS}^3 \mid \langle \tilde{x}_0, w \rangle \leq 2e^{2D} \quad \text{and} \quad \langle \tilde{\nu}_0, w \rangle \leq 2e^{2D}\}$$

which is a compact subset of AdS^3 , and the lemma follows. We now turn to the proof of the claim.

We fix n and consider the following functions:

$$\begin{aligned} a(t) &= -\langle \tilde{x}_0, \phi_n(\tilde{\zeta}(t)) \rangle, & a_\perp(t) &= -\langle \tilde{x}_0, \tilde{\nu}_n(\tilde{\zeta}(t)) \rangle, \\ a^*(t) &= -\langle \tilde{\nu}_0, \phi_n(\tilde{\zeta}(t)) \rangle, & a_\perp^*(t) &= -\langle \tilde{\nu}_0, \tilde{\nu}_n(\tilde{\zeta}(t)) \rangle. \end{aligned}$$

Notice that a is a positive function since \tilde{x}_0 and $\phi_n(\tilde{\zeta}(t))$ are contained in a spacelike surface. Moreover, since the surface \tilde{F}_n is convex, the plane orthogonal to $\tilde{\nu}_n(\tilde{\zeta}(t))$ is a support plane for \tilde{F}_n , so it is not difficult to check that also a_\perp is positive (see Figure 2).




FIGURE 2. The product $\langle \tilde{x}_0, \tilde{\nu}_n(\tilde{\zeta}(t)) \rangle$ is negative.

We can decompose \tilde{x}_0 as

$$\tilde{x}_0 = a(t)\tilde{\phi}_n(\tilde{\zeta}(t)) + a_\perp(t)\tilde{\nu}_n(\tilde{\zeta}(t)) + \tilde{v}(t)$$

with $\tilde{v}(t) \in T_{\phi_n(\tilde{\zeta}(t))}\tilde{F}_n$. Imposing $\langle \tilde{x}_0, \tilde{x}_0 \rangle = -1$, we deduce that $\|\tilde{v}(t)\| \leq a + a_\perp$.

On the other hand, $\|\tilde{v}(t)\| \leq \|\tilde{v}(t)\|_{\tilde{h}_n}$ and so

$$\begin{aligned} \dot{a} &= \langle \tilde{x}_0, \dot{\phi}_n \circ \tilde{\zeta} \rangle = \langle \tilde{v}, d\phi_n(\dot{\tilde{\zeta}}) \rangle \leq (a + a_\perp)\|\dot{\tilde{\zeta}}\|_{\tilde{h}_n} \\ \dot{a}_\perp &= \langle \tilde{x}_0, B_{\tilde{F}_n}(d\phi_n(\dot{\tilde{\zeta}})) \rangle \leq (a + a_\perp)\|B_{\tilde{F}_n}(\dot{\tilde{\zeta}})\|_{\tilde{h}_n} \leq (a + a_\perp)\frac{\theta_n}{2}\|\dot{\tilde{\zeta}}\|_{\tilde{h}_n^*}. \end{aligned}$$

Since $I_{S_n}^\#$ dominates both \tilde{h}_n and $\frac{\theta_n^2}{4}\tilde{h}_n^\star$ we see that

$$\begin{cases} (a + a_\perp)(0) = 2 \\ \dot{a} + \dot{a}_\perp \leq 2(a + a_\perp) \|\dot{\zeta}\|_{I_{S_n}^\#} \end{cases}$$

and by a simple integration we have $a + a_\perp \leq 2e^{2D}$.

A similar argument can be applied to a^\star and a_\perp^\star using the fact that the path $\tilde{\nu}_n(\zeta(t))$ is contained in the dual surface \tilde{F}_n^\star , that is the surface made of normal vectors of \tilde{F}_n . Indeed, there is a natural map $\phi_n^\star : \tilde{S} \rightarrow \tilde{F}_n^\star$ that sends a point \tilde{p} to the point dual to the plane tangent to \tilde{F}_n at $\phi_n(\tilde{p})$. The corresponding embedding data are

$$(37) \quad I_{\tilde{F}_n^\star} = \text{III}_{\tilde{F}_n} = \sin^2(\theta_n/2)h_n^\star, \quad B_{\tilde{F}_n^\star} = -1/\tan(\theta_n/2)b_n^{-1}.$$

In particular, F_n^\star is a past-convex spacelike surface, and the previous argument shows that $a^\star + a_\perp^\star \leq 2e^{2D}$. \square

Proof of Proposition 6.9. We will consider the product model of $\text{AdS}^3 = \mathbb{H}^2 \times S^1$, where the metric at some point $(\xi, e^{i\vartheta})$ is

$$g_{\text{AdS}^3} = g_{\mathbb{H}^2} - \chi(\xi)d\vartheta$$

where $\chi(\xi) = \cosh d_{\mathbb{H}^2}(\xi, \xi_0)$, where ξ_0 is some fixed point (see [7]).

By a lemma of Mess [20], the image of ϕ_n is the graph of some function $\mathbb{H}^2 \ni \xi \mapsto e^{is_n(\xi)} \in S^1$ that satisfies the following spacelike condition

$$(38) \quad \|\text{grad}(s_n)\| < 1/\chi.$$

We can also suppose that $\phi_n(\tilde{p}_0)$ is the point $\tilde{x}_0 = (\xi_0, 0)$ and the normal vector of \tilde{F}_n at \tilde{x}_0 is the unit vertical vector.

By (38), the functions s_n are uniformly Lipschitz on compact sets of \mathbb{H}^2 . So, up to subsequence, \tilde{F}_n converges to a surface \tilde{F}_∞ which is the graph of some limit function s_∞ , that verifies $\|\text{grad}(s_\infty)\| \leq 1/\chi$ almost every-where.

In order to show that \tilde{F}_∞ is spacelike, we need to prove that s_∞ verifies the strict inequality (38) almost everywhere. Notice that the projection $\pi_n : \tilde{F}_n \rightarrow \mathbb{H}^2$ increases the length, so the disk D in \mathbb{H}^2 with center $(\xi_0, 0)$ and radius r is contained in $\pi_n \circ \phi_n(B_{\tilde{F}_n}(\tilde{x}_0, r))$. By Lemma 6.11, the normal vectors of \tilde{F}_n on the cylinder based on D are contained in some compact subset K (independent of n).

Since the normal vector at $(\xi, s_n(\xi))$ is the vector

$$\frac{1}{\sqrt{1 - \chi^2 \|\text{grad}(s_n)\|^2}} \left(\text{grad}(s_n) + \frac{\partial}{\partial \vartheta} \right)$$

under the natural identification $T(\mathbb{H}^2 \times S^1) = T\mathbb{H}^2 \oplus TS^1$, we deduce that there exists ϵ depending on K , such that

$$\|\text{grad}(s_n)\| \leq (1 - \epsilon)/\chi$$

for every $\xi \in D$ and every n . This shows that \tilde{F}_∞ is spacelike.

Moreover, the restriction of the projection $\pi_n \circ \phi_n : (\tilde{S}, h) \rightarrow \mathbb{H}^2$ on $B_{\tilde{h}}(\tilde{p}_0, r)$ is C -Lipschitz, for some constant C depending only on r . Indeed, given a vector $\tilde{v} \in T_{\tilde{p}}\tilde{S}$, let $\tilde{v}_n = d\phi_n(\tilde{v})$ and $\tilde{u}_n = d\pi_n(\tilde{v}_n)$. We have that $\tilde{v}_n = \tilde{u}_n + \langle \text{grad}(s_n), \tilde{u}_n \rangle \frac{\partial}{\partial \vartheta}$, so $\cos^2(\theta_n/2)\tilde{h}_n(\tilde{v}, \tilde{v}) = \langle \tilde{v}_n, \tilde{v}_n \rangle \geq \|\tilde{u}_n\|^2 - \chi^2 \|\text{grad}(s_n)\|^2 \|\tilde{u}_n\|^2 \geq \epsilon \|\tilde{u}_n\|^2$.

Since $h_n \rightarrow h$, there is C' such that the identity map between (S, h) and (S, h_n) is C' -Lipschitz for every n . It follows, after taking a subsequence, $(\pi_n \circ \phi_n)$ converges to a map $\pi'_\infty : \tilde{S} \rightarrow \mathbb{H}^2$, so (ϕ_n) converges to the map $\phi_\infty(\tilde{p}) = (\pi'_\infty(\tilde{p}), s_\infty(\pi'_\infty(\tilde{p})))$. \square

We can prove now that the holonomy $\rho_n : \pi_1(S) \rightarrow \text{Isom}_0(\text{AdS}^3)$ of N_n converges to a limit representation ρ_∞ for which ϕ_∞ is equivariant.

Lemma 6.12. *If ϕ_n converges to a space-like embedding ϕ_∞ , then the representation ρ_n converges to a representation $\rho_\infty : \pi_1(S) \rightarrow \text{Isom}_0(\text{AdS}^3)$ such that \tilde{F}_∞ is ρ_∞ -equivariant.*

Moreover, the left and right components of ρ_∞ are discrete and faithful representations of $\pi_1(S)$ into $\text{PSL}_2(\mathbb{R})$.

Proof. First we prove that, for every $\gamma \in \pi_1(S)$, the sequence $\rho_n(\gamma)$ is bounded in $\text{Isom}_0(\text{AdS}^3)$.

Recall that we are assuming that $\phi_n(\tilde{p}_0) = \tilde{x}_0$ for all n and that the normal vectors $\tilde{\nu}_n(\tilde{p}_0)$ are equal to $\tilde{\nu}_0$. Now the $\rho_n(\gamma)(\tilde{x}_0) = \phi_n(\gamma\tilde{p}_0)$ form a sequence converging to $\bar{x}_0 = \phi_\infty(\gamma\tilde{p}_0)$ and the $\rho_n(\gamma)(\tilde{\nu}_0)$ converge to a unit timelike vector $\bar{\nu}_0$ at \bar{x}_0 orthogonal to some support plane of \tilde{F}_∞ .

This implies there is a bounded sequence of isometry of AdS^3 , says τ_n , such that

$$\tau_n \rho_n(\gamma)(\tilde{x}_0) = \tilde{x}_0, \quad \tau_n \rho_n(\gamma)(\tilde{\nu}_0) = \tilde{\nu}_0$$

Now the set of isometries that fix \tilde{x}_0 and $\tilde{\nu}_0$ is compact; so, after taking a subsequence, $\tau_n \rho_n(\gamma) \rightarrow \bar{\tau}$. Since up to passing to a subsequence we also have $\tau_n \rightarrow \tau_\infty$, we can deduce that $\rho_n(\gamma) \rightarrow \tau_\infty^{-1} \circ \bar{\tau}$.

To prove that ρ_n is converging, it is sufficient to check that two converging subsequences of ρ_n share the same limit. On the other hand, suppose that ρ_∞ is a limit of a subsequence of ρ_n , then ρ_∞ makes ϕ_∞ equivariant:

$$\phi_\infty(\gamma \tilde{p}) = \rho_\infty(\gamma) \phi_\infty(\tilde{p}) .$$

This relation uniquely determines the action of $\rho_\infty(\gamma)$ on \tilde{F}_∞ . Since two isometries of AdS^3 that coincide on a spacelike surface are equal the result follows.

The fact that the left and right representations $\pi_1(S) \rightarrow \text{PSL}_2(\mathbb{R})$ corresponding to ρ_∞ are faithful and discrete is a consequence of the fact that they are limit of faithful and discrete representations. \square

If (ϕ_{n_i}) is a convergent subsequence of (ϕ_n) , then Lemma 6.12 implies that N_{n_i} is a convergent sequence of spacetime. Let N_∞ be the limit of such spacetimes. Its holonomy is by definition the limit of the holonomies of the N_n . In particular, we can concretely realize N_n as an AdS metric g_{N_n} on $S \times \mathbb{R}$, in such a way that $g_{N_{n_i}}$ converges to an AdS metric g_{N_∞} as tensors on $S \times \mathbb{R}$ and $(S \times \mathbb{R}, g_{N_\infty}) \cong N_\infty$.

Proposition 6.13. *F_{n_i} converges to the lower boundary $\partial_- C(N_\infty)$ of the convex core of N_∞ . Moreover, the induced map*

$$\bar{\phi}_\infty : (S, h) \rightarrow \partial_- C(N_\infty)$$

is an isometry.

The proof of this proposition is based on the following lemma.

Lemma 6.14. *If N is a MGHC anti de Sitter spacetime and $N(k)$ is a Cauchy surface of constant curvature $k \leq -1$, then the Lorentzian distance of any point of $N(k)$ from the convex core of N is smaller than $\arctan \sqrt{|1+k|}$.*

Proof. We consider the point x_0 on $N(k)$ with the biggest distance from the convex core. If d is the distance between x_0 and the convex core, then it is well known that $d < \pi/2$ and there exists a timelike geodesic segment ς joining the point x_0 to a point y_0 on the boundary of the convex core with length equal to d [5].




FIGURE 3. Estimating the distance between $\tilde{N}(k)$ and $C(\tilde{N})$.

We consider now a lift $\tilde{\varsigma}$ of ς to the universal cover $\tilde{N} \subset \text{AdS}^3$ (see Figure 3). The plane Ξ through \tilde{y}_0 orthogonal to $\tilde{\varsigma}$ is a support plane for the lifting of the convex core. Let Ξ^d be the surface of points in AdS^3 whose distance from Ξ is d . It is a convex surface of constant curvature $-1/\cos^2(d)$. Clearly, \tilde{x}_0 lies on Ξ^d and $\tilde{N}(k)$ is contained in the convex side bounded by Ξ^d . In particular, $\tilde{N}(k)$ and Ξ^d are tangent at \tilde{x}_0 and, by the maximum principle, the curvature of $\tilde{N}(k)$ is less than the curvature of Ξ^d at \tilde{x}_0 .

We deduce that $k \leq -1/\cos^2(d)$, which implies that $\tan(d) \leq \sqrt{|1+k|}$. \square

Proof of Proposition 6.13. Since the metrics $g_{N_{n_i}}$ converge to g_{N_∞} , the convex cores of the N_{n_i} converge to the convex core of N_∞ . (Since the metrics converge, the holonomy representations converge and so that their limit set in $\partial_\infty \text{AdS}^3$ converge; therefore also their convex hulls, so the convex cores converge.)

In particular, the lower boundary $\partial_- C(N_{n_i})$ of the convex core of N_{n_i} converges to $\partial_- C(N_\infty)$. By Lemma 6.14, the distance of any point of F_{n_i} from $\partial_- C(N_{n_i})$ is smaller than $\theta_{n_i}/2$. This implies that F_{n_i} converges to $\partial_- C(N_\infty)$.

In order to prove that the map $\bar{\phi}_\infty : (S, h) \rightarrow \partial_- C(N_\infty)$ is an isometry, it is sufficient to show that $\bar{\phi}_\infty$ increases the distances. Indeed, both (S, h) and $\partial_- C(N_\infty)$ are hyperbolic surfaces and $\bar{\phi}_\infty$ is an homotopy equivalence.

We will prove that the lifting $\phi_\infty : (\tilde{S}, h) \rightarrow \partial_- C(\tilde{N}_\infty)$ increases the lengths. Given $\tilde{p}, \tilde{q} \in \tilde{S}$, we consider any path $\tilde{\zeta} : [0, 1] \rightarrow \tilde{S}$ connecting \tilde{p} and \tilde{q} such that

- $\partial_- C(\tilde{N}_\infty)$ is smooth at almost all points of $\tilde{\zeta}_\infty := \phi_\infty \circ \tilde{\zeta}$;
- $\ell(\tilde{\zeta}_\infty) \leq d_\infty(\tilde{x}_\infty, \tilde{y}_\infty) + \epsilon$.

where $\tilde{x}_\infty = \phi_\infty(\tilde{p})$ and $\tilde{y}_\infty = \phi_\infty(\tilde{q})$ and d_∞ is the distance in $\partial_- C(\tilde{N}_\infty)$.

In the model $\mathbb{H}^2 \times S^1$ of AdS^3 , the surfaces \tilde{F}_n are graphs of functions $e^{is_n} : \mathbb{H}^2 \rightarrow S^1$ converging to $e^{is_\infty} : \mathbb{H}^2 \rightarrow S^1$ such that $\partial_- C(\tilde{N}_\infty)$ is the graph of e^{is_∞} .

We have $\tilde{\zeta}_\infty(t) = (\xi(t), e^{is_\infty(\xi(t))})$ with $\xi : [0, 1] \rightarrow \mathbb{H}^2$ Lipschitz function. Take $\tilde{\zeta}_n(t) = (\xi(t), e^{is_n(\xi(t))})$. For any smooth point $\tilde{x} = (\xi, e^{is_\infty(\xi)})$ of $\partial_- C(\tilde{N}_\infty)$ we have $\text{grad}(s_n)(\xi) \rightarrow \text{grad}(s_\infty)(\xi)$. Indeed, by convexity, tangent planes of \tilde{F}_n converge to support planes of $\partial_- C(\tilde{N}_\infty)$.

By Lebesgue Theorem we have

$$\ell(\tilde{\zeta}_n) = \int_0^1 \sqrt{\|\dot{\xi}\|^2 - \chi(\xi) \langle \dot{\xi}, \text{grad}(s_n) \rangle^2} dt \rightarrow \int_0^1 \sqrt{\|\dot{\xi}\|^2 - \chi(\xi) \langle \dot{\xi}, \text{grad}(s_\infty) \rangle^2} dt = \ell(\tilde{\zeta}_\infty)$$

since $\sqrt{\|\dot{\xi}\|^2 - \chi(\xi) \langle \dot{\xi}, \text{grad}(s_n) \rangle^2}$ are all dominated by $\|\dot{\xi}\|$, which is an integrable function.

Since the endpoints of $\tilde{\zeta}_n$ correspond to points $\tilde{x}_n = \varphi_n(\tilde{p}_n)$ and $\tilde{y}_n = \varphi_n(\tilde{q}_n)$ with $\tilde{p}_n \rightarrow \tilde{p}$ and $\tilde{q}_n \rightarrow \tilde{q}$, we deduce that $d_{\tilde{h}}(\tilde{p}, \tilde{q}) \leq d_\infty(\tilde{x}_\infty, \tilde{y}_\infty) + \epsilon$. Since ϵ can be chosen arbitrarily small,

$$(39) \quad d_{\tilde{h}}(\tilde{p}, \tilde{q}) \leq d_\infty(\tilde{x}_\infty, \tilde{y}_\infty).$$

□

So far, we have shown that N_n is contained in a compact subset of the space of MGH AdS structures, and any convergent subsequence of $\phi_n : S \rightarrow N_n$ converges to an isometric embedding $\bar{\phi}_\infty : (S, h) \rightarrow N_\infty$, whose image is the lower boundary $\partial_- C(N_\infty)$ of the convex core of N_∞ .

Let λ_∞ be the bending lamination of $\partial_- C(N_\infty)$. We will prove that $\lambda_\infty = \lambda/2$. Since the length spectrum of the third fundamental form \mathbb{I}_{F_n} converges to the intersection spectrum of $\lambda/2$, it is sufficient to prove that it converges also to the intersection spectrum of the bending lamination of $\partial_- C(N_\infty)$.

Now, let \tilde{F}_n^* be the surface dual to \tilde{F}_n . Points of \tilde{F}_n^* are dual to tangent planes of \tilde{F}_n and \tilde{F}_n^* is a past-convex surface of constant curvature $-1/\sin^2(\theta_n/2)$ as (37) shows.

Clearly, \tilde{F}_n^* is invariant under the holonomy action of $\pi_1(S)$, so it is contained in \tilde{N}_n^* and its quotient is a Cauchy surface F_n^* of \tilde{N}_n^* . By (37), the length spectrum of \tilde{F}_n^* is equal to the length spectrum of the third fundamental form \mathbb{I}_{F_n} .

The boundary of the domain \tilde{N}_∞ in AdS^3 is the union of two disjoint achronal surfaces: the past and the future singularities of \tilde{N}_∞ , that are clearly invariant under the action of $\pi_1(S)$.

By Proposition 6.7, the length spectrum of F_n^* converges to the length spectrum of the action of $\pi_1(S)$ on the future singularity of \tilde{N}_∞^* (notice indeed that since F_n^* is past-convex, in order to apply Proposition 6.7 we need to exchange the time orientation). On the other hand, by [5] the length spectrum of the future singularity of \tilde{N}_∞^* coincides with the intersection spectrum of the bending lamination of the lower boundary of the convex core of N_∞ .

Combining these facts, we deduce that

$$\ell_{\mathbb{I}_{F_n}}(\gamma) \rightarrow \iota(\lambda_\infty, \gamma)$$

so $\lambda_\infty = \lambda/2$.

6.3. Asymptotic behavior of the measures $\text{tr}(b_n)\omega_{h_n}$. Let $(h_n, h_n^*)_{n \in \mathbb{N}}$ be a sequence of normalized hyperbolic metrics on S such that h_n converges to h and h_n^* converges to $[\lambda]$ in Thurston boundary of Teichmüller space, and denote by b_n the operator associated to (h_n, h_n^*) provided by Corollary 1.5.

Moreover, let $(\theta_n)_{n \in \mathbb{N}}$ be a sequence such that $\theta_n \ell_{h_n^*}$ converges to $\iota(\lambda, \bullet)$ in the sense of spectra of closed curves.

In this section we study the asymptotic behavior of $\text{tr}(b_n)$: roughly speaking, it concentrates around the h -geodesic representative of λ . Hence, we will always refer to λ as to such an h -geodesic representative.

These results will turn useful in the proof of Theorem 6.1 and in Section 7.

Proposition 6.15. *Let $V \subset S$ a closed subsurface such that ∂V is smooth and does not intersect λ . Call λ_V the h -geodesic sublamination $\lambda \cap V$. Then*

$$\theta_n \int_V \text{tr}(b_n) \omega_{h_n} \rightarrow \ell_h(\lambda_V).$$

where ω_g is the area form associated to g .

In order to prove Proposition 6.15, we need the following lemma that is analogous to Lemma 6.14.

Lemma 6.16. *Let M be a hyperbolic end associated to some projective structure on S , and let $M(k)$ be the surface of constant curvature k with $k \in [-1, 0)$. Then the distance of any point of $M(k)$ from the boundary of M is at most $\text{arctanh} \sqrt{1+k}$.*

The proof of Lemma 6.16 is essentially the same as in Lemma 6.14. We leave the details to the reader.

Corollary 6.17. *The family of isometric immersions $\sigma_n : (\tilde{S}, \cosh^2(\theta_n/2)\tilde{h}) \rightarrow \mathbb{H}^3$ converges to a bending map $\sigma_\infty : (\tilde{S}, h) \rightarrow \mathbb{H}^3$, with bending lamination $\lambda/2$.*

Proof. Since σ_n are uniformly Lipschitz as maps $(\tilde{S}, \tilde{h}) \rightarrow \mathbb{H}^3$, they converge up to subsequences to a locally convex surface. Combining Proposition 6.2 and Lemma 6.16, we deduce that this surface is the bent surface corresponding to $Gr_{\lambda/2}(S)$. \square

Proof of Proposition 6.15. We consider the embedding $\bar{\sigma}_n : S \rightarrow S_n \in M_n$ inside a hyperbolic end M_n such that

$$I_{S_n} = \cosh^2(\theta_n/2)h_n \quad B_{S_n} = \tanh(\theta_n/2)b_n.$$

The projective structure G_n at the ideal boundary of M_n converges to $G_\infty = Gr_{\lambda/2}(h)$ (Proposition 6.2). Call M_∞ the hyperbolic end determined by the \mathbb{CP}^1 -surface G_∞ .

A simple computation shows that the area element of S_n with respect to $I_{S_n}^\#$ is $\omega_{I_{S_n}^\#} = (\cosh^2(\theta_n/2) + \sinh^2(\theta_n/2) + \sinh(\theta_n/2) \cosh(\theta_n/2) \text{tr}(b_n)) \omega_{h_n}$ and so the area of $\bar{\sigma}_n(V)$ is

$$(40) \quad \text{Area}_{I_{S_n}^\#}(V) = \left(\text{Area}_{h_n}(V) + \frac{\theta_n}{2} \int_V \text{tr}(b_n) \omega_{h_n} \right) (1 + o(\theta_n))$$

As before, we can identify $M_n \cup G_n \cong (S \times [0, \infty), g_{M_n}) \cup S \times \{\infty\}$ in such a way that:

- the developing map $dev_n : \tilde{S} \times [0, \infty) \rightarrow \mathbb{H}^3$ converges to dev_∞ (and so $g_{M_n} \rightarrow g_{M_\infty}$);
- $\bar{\sigma}_n$ converges to the pleated surface $\bar{\sigma}_\infty : S \rightarrow (S \times \{0\}, g_{M_\infty})$

Call $\partial^d M_n$ the surface in M_n at distance d from the boundary and let $\Pi_{\partial^d M_n} : G_n \rightarrow \partial^d M_n$ the projection introduced in Lemma 6.10.

There exists two numbers $\epsilon_n < \delta_n$ such that S_n is contained between $\partial^{\epsilon_n} M_n$ and $\partial^{\delta_n} M_n$ and, by Lemma 6.16, $\delta_n \rightarrow 0$ as $n \rightarrow +\infty$.

By the monotonicity result proved in [28],

$$\Pi_{\partial^{\epsilon_n} M_n}^*(I_{\partial^{\epsilon_n} M_n}^\#) \leq \Pi_{S_n}^*(I_{S_n}^\#) \leq \Pi_{\partial^{\delta_n} M_n}^*(I_{\partial^{\delta_n} M_n}^\#).$$

If λ_n is the bending lamination of M_n , the grafted metric on $\partial^d M_n$ makes it isometric to $e^{2d} g_{G_n}$, where g_{G_n} is Thurston metric on the projective surface $G_n = Gr_{\lambda_n}(\partial M_n, g_{M_n})$.

So we deduce that

$$e^{2\epsilon_n} \omega_{G_n} \leq \Pi_{S_n}^*(\omega_{I_{S_n}^\#}) \leq e^{2\delta_n} \omega_{G_n}$$

and so

$$(41) \quad e^{2\epsilon_n} \text{Area}_{G_n}(\Pi_{S_n}^{-1}(\bar{\sigma}_n(V))) \leq \text{Area}_{I_{S_n}^\#}(V) \leq e^{2\delta_n} \text{Area}_{G_n}(\Pi_{S_n}^{-1}(\bar{\sigma}_n(V))).$$

Since $G_n \rightarrow G_\infty$, their Thurston metrics converge to g_{G_∞} . We claim that $\Pi_{S_n}^{-1}(\bar{\sigma}_n(V))$ converge to $\Pi_{S_\infty}^{-1}(\bar{\sigma}_\infty(V))$ in the Hausdorff sense, and so

$$(42) \quad \text{Area}_{I_{S_n}^\#}(V) \rightarrow \text{Area}_{G_\infty}(\Pi_{S_\infty}^{-1}(\bar{\sigma}_\infty(V))) = \text{Area}_h(V) + \frac{1}{2} \ell_h(\lambda_V)$$

by Equation (41).

The result will follow by comparing Equations (40) and (42).

In order to prove the claim, it is enough to prove that $\partial \Pi_{S_n}^{-1}(\bar{\sigma}_n(V)) \rightarrow \partial \Pi_{S_\infty}^{-1}(\bar{\sigma}_\infty(V))$, which would follow from the fact that $\Pi_{S_n}^{-1} \circ \bar{\sigma}_n|_{\partial V}$ converges to $\Pi_{S_\infty}^{-1} \circ \bar{\sigma}_\infty|_{\partial V}$.

Notice that Π_{S_n} is a diffeomorphism, and $\Pi_{S_\infty}^{-1} \circ \bar{\sigma}_\infty|_{\partial V}$ is well-defined and continuous since ∂V does not intersect λ .

Let $(p_n)_{n \in \mathbb{N}}$ be a sequence of points in ∂V such that $p_n \rightarrow p$. The point $\Pi_{S_n}^{-1}(\bar{\sigma}_n(p_n))$ is the ideal point of the horocycle U_n tangent to S_n at $\bar{\sigma}_n(p_n)$. By convexity of the surfaces S_n , one can easily see that U_n converges to a horocycle U_∞ tangent to S_∞ at $\bar{\sigma}_\infty(p)$. Since $p \notin \lambda$, such a U_∞ is unique and so its ideal point is necessarily $\Pi_{S_\infty}^{-1}(p)$. \square

Corollary 6.18. *Let h_n be a sequence of hyperbolic metrics converging to h , and let h_n^* be a diverging sequence of metrics. Then, $(\theta_n \ell_{h_n^*}(\gamma))_{n \in \mathbb{N}}$ is bounded for every $\gamma \in \pi_1(S)$ if and only if $\theta_n \int_S \text{tr}(b_n) \omega_{h_n}$ is bounded.*

6.4. Proof of Theorems 1.12 and 6.1.

Proof of Theorem 1.12. Let us fix a hyperbolic metric h on S and a sequence of hyperbolic metrics h_n^* converging to a point $[\lambda]$ in Thurston boundary of \mathcal{T} . Fixing a sequence θ_n such that $\theta_n \ell_{h_n^*}$ converges to $\iota(\lambda, \bullet)$, let us set

$$L_{e^{i\theta_n}}(h, h_n^*) = (h_n^1, h_n^2).$$

In subsection 6.2 we have shown that $h_n^1 \rightarrow E_{\lambda/2}(h)$. To conclude the proof we need to prove that

$$(43) \quad \theta_n \ell_{h_n^2} \rightarrow \iota(\lambda, \bullet),$$

The main issue to prove (43) is to show that for every $\gamma \in \pi_1(S)$, there is $C = C(\gamma)$ such that

$$(44) \quad \theta_n \ell_{h_n^2}(\gamma) < C.$$

Corollary 6.18 indicates that, in order to prove (44), it is sufficient to bound

$$\theta_n \int_S \text{tr}(b'_n) \omega_{h_n^1}$$

where b'_n is the h_n^1 -self-adjoint operator such that $h_n^2 = h_n^1(b'_n \bullet, b'_n \bullet)$. Now we have that $b'_n = \beta'_n b_n \beta_n^{-1}$ where $\beta_n = \cos(\theta_n/2)E + \sin(\theta_n/2)Jb_n$ and $\omega_{h_n^1} = \det(\beta_n) \omega_h = \omega_h$. So, $\theta_n \int_S \text{tr}(b'_n) \omega_{h_n^1} = \theta_n \int_S \text{tr}(b_n) \omega_h$ that in turn is bounded since $\theta_n \ell_{h_n^*}(\gamma)$ is bounded for every $\gamma \in \pi_1(S)$ by hypothesis.

It follows that there exists a measured geodesic lamination μ such that, up to passing to a subsequence, $\theta_n \ell_{h_n^2} \rightarrow \iota(\mu, \bullet)$. To show that $\mu = \lambda$, notice that by Proposition 6.8 we have that

$$(45) \quad L_{e^{i\theta_n}}^1(h_n^1, h_n^2) \rightarrow E_{\mu/2}(h_\infty)$$

On the other hand, we have that

$$L_{e^{i\theta_n}}^1(h_n^1, h_n^2) = L_{e^{i\theta_n}}^1(L_{e^{i\theta_n}}(h, h_n^*)) = L_{e^{2i\theta_n}}^1(h, h_n^*).$$

So, applying again Proposition 6.8, we obtain that

$$(46) \quad L_{e^{i\theta_n}}^1(h_n^1, h_n^2) \rightarrow E_\lambda(h) = E_{\lambda/2}(h_\infty).$$

Comparing (45) and (46) we conclude that $E_{\lambda/2}(h_\infty) = E_{\mu/2}(h_\infty)$ and so $\lambda = \mu$. \square

Proof of Theorem 6.1. Let h and h_n as above. For all $z = t + is \in \overline{\mathbb{H}}$, we have to prove

$$(47) \quad P'_{\theta_n z}(h, h_n^*) \rightarrow Gr_{s\lambda/2}(E_{-t\lambda/2}(h)).$$

Recall that $P'_{\theta_n z}(h, h_n^*) = SGr_{\theta_n s} \circ L'_{-\theta_n t}(h, h_n^*)$. Note that if we put

$$L'_{-\theta_n t}(h, h_n^*) = (h_n^1, h_n^2)$$

Theorem 1.12 shows that $h_n^1 \rightarrow h_\infty = E_{-t\lambda/2}(h)$ and $\theta_n \ell_{h_n^2} \rightarrow \lambda$.

Applying Proposition 6.2 we conclude that

$$P'_{\theta_n z}(h, h_n^*) = SGr_{\theta_n s}(h_n^1, h_n^2) \rightarrow Gr_{s\lambda/2}(E_{-t\lambda/2}(h)).$$

\square

6.5. Convergence of the distances. The aim of this section is to study the asymptotic behavior of the sequence of distances induced by the metrics $\theta_n^2 h_n^*$ introduced in the previous section. By our assumption, the length spectrum of h_n^* rescaled by θ_n converges to the intersection with λ . Notice that this assumption only concerns the isotopy class of h_n^* . On the other hand, once we concretely fix h , the metric h_n^* is uniquely determined in its isotopy class by requiring the identity map $(S, h) \rightarrow (S, h_n^*)$ is minimal Lagrangian. The result we consider in this section deals with the asymptotic behavior of h_n^* considered as concrete metrics on S . Clearly, these results are valid for this choice of gauge, and are no longer valid if we change h_n^* by some isotopy.

Notice that the representative λ of a point in Thurston boundary of $\mathcal{T}(S)$ can be chosen to be a measured geodesic lamination for any hyperbolic metric on S . In order to study the behavior of h_n^* , it is natural to fix λ as the concrete measured geodesic realization with respect to the metric h .

Let us fix a universal cover $\tilde{S} \rightarrow S$ and denote by \tilde{h} and \tilde{h}_n^* the pull-back of the metrics h and h_n^* on \tilde{S} . Finally let $\tilde{\lambda}$ be the pull-back of λ on \tilde{S} .

Proposition 6.19. *For every $\tilde{p}, \tilde{q} \in \tilde{S} \setminus \tilde{\lambda}$ we have*

$$(48) \quad \theta_n d_{\tilde{h}_n^*}(\tilde{p}, \tilde{q}) \rightarrow \iota(\tilde{\alpha}, \tilde{\lambda})$$

where $\tilde{\alpha}$ is any smooth path in \tilde{S} joining \tilde{p} to \tilde{q} and meeting each leaf of $\tilde{\lambda}$ at most once and transversely. Moreover, the convergence is uniform on compact subsets of $\tilde{S} \setminus \tilde{\lambda}$.

Theorem 1.13 is a direct consequence of this statement, so that its proof will be a consequence of the proof of this proposition.

First, notice that it is sufficient to prove Proposition 6.19 after rescaling θ_n and λ by some arbitrary factor. In particular, we may assume that the projective surface $Gr_{\lambda/2}(S, h)$ is quasi-Fuchsian. This technical assumption will simplify some steps of the proof.

First we show that

Lemma 6.20.

$$(49) \quad \liminf \theta_n d_{\tilde{h}_n^*}(\tilde{p}, \tilde{q}) \geq \iota(\tilde{\alpha}, \tilde{\lambda}).$$

Proof. Let M_n be the hyperbolic end introduced in Section 6.1 with $h_n = h$ and let $\sigma_n : \tilde{S} \rightarrow \tilde{M}_n$ be the lifting of the embedding $\bar{\sigma}_n : S \rightarrow M_n$. By Proposition 6.2, M_n converges to the hyperbolic end facing $Gr_{\lambda/2}(S, h)$. In particular, we may assume that M_n are all quasi-Fuchsian, so that \tilde{M}_n is a concave region of \mathbb{H}^3 .

By Corollary 6.17, the family of embeddings $(\sigma_n)_{n \in \mathbb{N}}$ converges to the bending map $\sigma_\infty : (\tilde{S}, \tilde{h}) \rightarrow \mathbb{H}^3$, with bending lamination $\tilde{\lambda}/2$.

Let $\tilde{r}_n^* : \tilde{S}_n^* \rightarrow \partial \tilde{M}_n^*$ be the 1-Lipschitz map defined in Lemma 6.5. We denote by $\Xi_n(\tilde{p})$ the plane in \mathbb{H}^3 corresponding to $\tilde{r}_n^*(\sigma_n^*(\tilde{p}))$ and by $\Xi_n(\tilde{q})$ the plane corresponding to $\tilde{r}_n^*(\sigma_n^*(\tilde{q}))$.

We have that $\Xi_n(\tilde{p})$ and $\Xi_n(\tilde{q})$ are both support planes of $\partial \tilde{M}_n$. Let us put $\iota_n = d_{\partial \tilde{M}_n^*}(\tilde{r}_n^*(\sigma_n^*(\tilde{p})), \tilde{r}_n^*(\sigma_n^*(\tilde{q})))$.

Since \tilde{r}_n decreases the distances we deduce that

$$(50) \quad \iota_n \leq d_{\mathbb{H}^3}(\sigma_n(\tilde{p}), \sigma_n(\tilde{q})) \sim \frac{\theta_n}{2} d_{\tilde{h}_n^*}(\tilde{p}, \tilde{q}).$$

We claim that $\Xi_n(\tilde{p})$ and $\Xi_n(\tilde{q})$ converge to the support planes of $\partial \tilde{M}_\infty$ at $\sigma_\infty(\tilde{p})$ and $\sigma_\infty(\tilde{q})$ respectively.

Let us explain how the conclusion follows. The claim ensures that it is possible to construct a sequence of arcs $\tilde{\alpha}_n : [0, 1] \rightarrow \partial \tilde{M}_n$ such that

- each $\tilde{\alpha}_n$ intersects each leaf of $\tilde{\lambda}_n$ at most once and transversely; moreover end-points of $\tilde{\alpha}_n$ are contained in $\partial \tilde{M}_n \setminus \tilde{\lambda}_n$;
- end-points of $\tilde{\alpha}_n$ are close to $\Xi_n(\tilde{p})$ and $\Xi_n(\tilde{q})$ in the sense that $d_{\partial \tilde{M}_n^*}(\tilde{\alpha}_n^*(0), \tilde{r}_n^*(\sigma_n^*(\tilde{p})))$ and $d_{\partial \tilde{M}_n^*}(\tilde{\alpha}_n^*(1), \tilde{r}_n^*(\sigma_n^*(\tilde{q})))$ converge to 0.
- $\tilde{\alpha}_n$ converges to a path $\tilde{\alpha}_\infty$ connecting the stratum of $\partial \tilde{M}_\infty \setminus \tilde{\lambda}$ containing $\sigma_\infty(\tilde{p})$ with the stratum containing $\sigma_\infty(\tilde{q})$

Thus we have $\iota(\tilde{\lambda}/2, \tilde{\alpha}) = \iota(\tilde{\lambda}/2, \tilde{\alpha}_\infty) = \lim_n \iota(\tilde{\lambda}_n, \tilde{\alpha}_n)$. By (32) $\iota(\tilde{\lambda}_n, \tilde{\alpha}_n) = d_{\partial \tilde{M}_n^*}(\tilde{\alpha}_n^*(0), \tilde{\alpha}_n^*(1))$, so we conclude that

$$\iota(\tilde{\lambda}/2, \tilde{\alpha}) = \lim_n \iota_n$$

and the conclusion of the Lemma follows from (50).

In order to prove the claim recall that \tilde{r}_n^* sends any point \tilde{x}^* of \tilde{S}_n^* to a point in $\partial \tilde{M}_n^*$ that lies in the past of \tilde{x}^* (Lemma 6.5).




FIGURE 4. Convergence of the support planes.

This implies that $\Xi_n(\tilde{p})$ and $T_{\sigma_n(\tilde{p})}\tilde{S}_n$ are disjoint planes. In particular we deduce that $\Xi_n(\tilde{p})$ separates $\sigma_n(\tilde{p})$ from $\partial \tilde{M}_n$ (see Figure 4). This easily implies that $\Xi_n(\tilde{p})$ converges to the support plane of $\partial \tilde{M}_\infty$ at $\sigma_\infty(\tilde{p})$ (that is unique by our assumption that \tilde{p} does not lie on $\tilde{\lambda}$). Analogously $\Xi_n(\tilde{q})$ converges to the support plane of $\partial \tilde{M}_\infty$ at $\sigma_\infty(\tilde{q})$. \square

To conclude the proof of Proposition 6.19 we need to show that

$$(51) \quad \limsup \theta_n d_{\tilde{h}_n^*}(\tilde{p}, \tilde{q}) \leq \iota(\tilde{\alpha}, \tilde{\lambda}).$$

In order to estimate $\theta_n d_{\tilde{h}_n^*}(\tilde{p}, \tilde{q})$, we need the following result.

Lemma 6.21. *Let \tilde{U} be any convex smooth surface in \mathbb{H}^3 and let $\tilde{x}, \tilde{y} \in \tilde{U}$ such that the support planes $\Xi_{\tilde{x}}$ and $\Xi_{\tilde{y}}$ at \tilde{x} and \tilde{y} intersect. Then the distance between \tilde{x} and \tilde{y} with respect to the third fundamental form of \tilde{U} is less than the angle between $\Xi_{\tilde{x}}$ and $\Xi_{\tilde{y}}$.*

We first prove a 2-dimensional version of this lemma.

Sublemma 6.22. *Let $\tilde{\zeta}$ be a convex curve in \mathbb{H}^2 joining two points \tilde{x}, \tilde{y} . Suppose that the support lines $\tilde{l}_{\tilde{x}}$ and $\tilde{l}_{\tilde{y}}$ at \tilde{x} and \tilde{y} intersect. Then the angle they form is bigger than the integral of the curvature of $\tilde{\zeta}$.*

Proof. By Gauss-Bonnet formula, the area bounded by $\tilde{l}_{\tilde{x}}$, $\tilde{l}_{\tilde{y}}$ and $\tilde{\zeta}$ is equal to the difference between the wanted angle and the integral of the curvature. \square

Proof of Lemma 6.21. First we translate the condition that planes intersect in terms of a condition of dual points \tilde{x}^* and \tilde{y}^* . Recalling that \tilde{x}^* and \tilde{y}^* are unit vector orthogonal to the planes (and pointing in the concave region bounded by \tilde{U}), we easily derive that $\Xi_{\tilde{x}}$ and $\Xi_{\tilde{y}}$ intersect if and only if the segment $\tilde{\kappa}^*$ joining \tilde{x}^* to \tilde{y}^* in $d\mathbb{S}^3$ is spacelike, in which case the angle between $\Xi_{\tilde{x}}$ and $\Xi_{\tilde{y}}$ coincides with the length of $\tilde{\kappa}^*$.

Let Π be a timelike plane containing $\tilde{\kappa}^*$. Since \tilde{U}^* is an achronal surface, $\tilde{U}^* \cap \Pi$ is a curve containing \tilde{x}^* and \tilde{y}^* . Let $\tilde{\zeta}^*$ be the segment on $\tilde{U}^* \cap \Pi$ connecting \tilde{x}^* to \tilde{y}^* . Clearly, the length of $\tilde{\zeta}^*$ is greater than the distance on \tilde{U}^* between \tilde{x}^* and \tilde{y}^* . In order to conclude, it is sufficient to prove that the length of $\tilde{\zeta}^*$ is less than the length of $\tilde{\kappa}^*$.

Notice that this is a 2-dimensional problem. In fact, let Υ be the timelike linear 3-space in $\mathbb{R}^{3,1}$ such that $\Pi = \Upsilon \cap d\mathbb{S}^3$. The intersection $\Upsilon \cap \mathbb{H}^3$ is a hyperbolic plane denoted by Λ . Points on Π correspond to planes of \mathbb{H}^3 that orthogonally meet Λ . In particular, points on Π bijectively correspond to lines on Λ and points of Λ correspond to spacelike lines of Π .

Consider the curve $\tilde{\zeta}$ on Λ of points corresponding to support lines of $\tilde{\zeta}^*$. Notice that support lines at the endpoints of $\tilde{\zeta}$ are $\Xi_{\tilde{x}} \cap \Lambda$ and $\Xi_{\tilde{y}} \cap \Lambda$, so the angles these lines form is equal to the angle between $\Xi_{\tilde{x}}$ and $\Xi_{\tilde{y}}$ and it is equal to the length of $\tilde{\kappa}^*$. On the other hand the length of $\tilde{\zeta}^*$ is equal to the integral of the curvature of $\tilde{\zeta}$. Thus the conclusion follows from Sublemma 6.22. \square

Given any geodesic $\tilde{\zeta} \in (\tilde{S}, \tilde{h})$, we say that $\sigma_n(\tilde{\zeta})$ is a short path if the support planes at $\sigma_n(\tilde{\zeta}(0))$ and $\sigma_n(\tilde{\zeta}(1))$ intersect. If $\sigma_n(\tilde{\zeta})$ is a short path, then we denote by $\eta_n(\tilde{\zeta}) \in (0, \pi)$ the angle between the support planes at its endpoints.

Analogously, we say that $\sigma_\infty(\tilde{\zeta})$ is a short path if the endpoints of $\sigma_\infty(\tilde{\zeta})$ are outside the bending lamination and the corresponding support planes intersect. In this case, $\eta_\infty(\tilde{\zeta})$ is the angle between such planes.

Clearly, if $\sigma_\infty(\tilde{\zeta})$ is a short path, then $\sigma_n(\tilde{\zeta})$ is definitively a short path and $\eta_n(\tilde{\zeta}) \rightarrow \eta_\infty(\tilde{\zeta})$.

Lemma 6.23. *There exists ϵ_0 such that if $\tilde{\alpha} : [0, 1] \rightarrow \tilde{S}$ is a geodesic path for \tilde{h} of length less than ϵ_0 , then*


- $\sigma_\infty(\tilde{\alpha})$ is a short path;
- $\iota(\tilde{\alpha}, \tilde{\lambda}) \leq \eta_\infty(\tilde{\alpha}) \leq (1 + \ell_{\tilde{h}}(\tilde{\alpha}))\iota(\tilde{\alpha}, \tilde{\lambda})$

Proof. The first point easily follows, since \tilde{S}_∞ is invariant under the action of a co-compact group of isometries of \mathbb{H}^3 .

About the second point, notice that the first inequality is given by Lemma 6.21. The second inequality is more subtle. Choosing ϵ_0 sufficiently small, we can suppose that either $\tilde{\alpha}$ intersects only one isolated leaf of $\tilde{\lambda}$ or it intersects no isolated leaf. In the first case the second inequality is obvious.

Up to taking a smaller ϵ_0 , we may suppose that, if T is a hyperbolic triangle with an edge e of length $l \leq \epsilon_0$ and the angles ϑ_1, ϑ_2 adjacent to e less than $\pi/4$, then the area of T is less than $l\vartheta_2$. Now take a geodesic $\tilde{\alpha}$ on \tilde{S} of length less than ϵ which does not intersect the isolated leaves of the lamination. Taking any subdivision $\tilde{\alpha}(0) = \tilde{p}_1, \dots, \tilde{p}_{m+1} = \tilde{\alpha}(1)$ of $\tilde{\alpha}$, we consider the support planes Ξ_i of $\sigma_\infty(\tilde{S})$ at $\sigma_\infty(\tilde{p}_i)$. If the subdivision is sufficiently fine, then the angles ϑ_i between Ξ_i and Ξ_{i+1} are less than $\pi/4$. Now consider the boundary ∂K of the convex set K obtained by intersecting the half-spaces bounded by Ξ_1, \dots, Ξ_{m+1} and containing $\sigma_\infty(\tilde{S})$. Notice that ∂K is a finite bent surface: indeed, its bending lines are $\Xi_i \cap \Xi_{i+1}$ for every i such that Ξ_i and Ξ_{i+1} are different. We claim that $\eta_\infty(\tilde{\alpha}) \leq (1 + \ell_{\tilde{h}}(\tilde{\alpha})) \sum \vartheta_i$. Taking a sequence of arbitrary fine subdivisions, we have that $\sum \vartheta_i \rightarrow \iota(\tilde{\alpha}, \tilde{\lambda})$, so the second inequality follows from the claim.

In order to prove the claim we use an inductive argument. Notice that, if Ξ_1, Ξ_2, Ξ_3 are distinct, then $\Xi_1 \cap \Xi_2 \cap \Xi_3 = \emptyset$. Thus, there is a plane Λ orthogonal to all of them. The triangle T obtained by intersecting $\Lambda \cap \Xi_1$, $\Lambda \cap \Xi_2$ and $\Lambda \cap \Xi_3$ has angles ϑ_1, ϑ_2 and $\pi - \bar{\vartheta}_1$, where $\bar{\vartheta}_1$ is the angle formed by Ξ_1 and Ξ_3 .



Notice that the length of the edge between ϑ_1 and ϑ_2 is less than the distance between \tilde{p}_1 and \tilde{p}_2 , and so it is smaller than ϵ_0 . We conclude that the area of T is less than $d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_2)\vartheta_2$. In particular, we deduce that $\bar{\vartheta}_1 \leq \vartheta_1 + \vartheta_2(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_2))$.

If at least two planes among Ξ_1, Ξ_2, Ξ_3 coincide, then the area of T is zero, and so $\bar{\vartheta}_1 = \vartheta_1 + \vartheta_2 \leq \vartheta_1 + \vartheta_2(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_2))$.

Apply now the same argument to the planes Ξ_1, Ξ_3, Ξ_4 . If $\bar{\vartheta}_2$ is the angle formed by Ξ_1 and Ξ_4 , then $\bar{\vartheta}_2 \leq \vartheta_1 + \vartheta_3(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_3)) \leq \bar{\vartheta}_1 + \vartheta_2(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_2)) + \vartheta_3(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_3))$. Iterating this procedure, we deduce that the angle between Ξ_1 and Ξ_m is bounded by $\vartheta_1 + \vartheta_2(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_2)) + \vartheta_3(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_3)) + \dots + \vartheta_m(1 + d_{\tilde{h}}(\tilde{p}_1, \tilde{p}_m))$ and this quantity is less than $(1 + \ell_{\tilde{h}}(\tilde{\alpha})) \sum \vartheta_i$. \square

We can now prove (51). Fix $\epsilon < \epsilon_0$ and subdivide the geodesic $\tilde{\alpha}$ joining \tilde{p} to \tilde{q} into segments $\tilde{\alpha}_i$ of length less than ϵ . Let $\tilde{p} = \tilde{p}_1, \tilde{p}_2, \dots, \tilde{p}_{m+1} = \tilde{q}$ be endpoints of such subdivision.

For n large, $\sigma_n(\tilde{\alpha}_i)$ are short paths and so by Lemma 6.21

$$\limsup d_{\mathbb{H}_{\tilde{S}_n}}(\tilde{p}, \tilde{q}) \leq \sum_{i=1}^m \eta(\tilde{\alpha}_i)$$

On the other hand, applying Lemma 6.23 we deduce that

$$\sum_{i=1}^m \eta(\tilde{\alpha}_i) \leq \iota(\tilde{\alpha}, \tilde{\lambda})(1 + \epsilon)$$

The uniform convergence follows from the fact that the whole argument works as well, if we consider sequences of points $\tilde{p}_n \rightarrow \tilde{p}$ and $\tilde{q}_n \rightarrow \tilde{q}$ belonging to a compact subset of $\tilde{S} \setminus \tilde{\lambda}$. This concludes the proof of Proposition 6.19.

7. BEHAVIOR OF THE CENTERS

In this section we want to discuss the behavior of the centers c_n when h is fixed and $h_n^* \rightarrow [\lambda]$ in Thurston compactification. Our aim is to prove that the limit point(s) of c_n does not only depend on h and $[\lambda]$ but also on the sequence h_n^* . As a consequence, we will see that the analog of Theorem 1.12 does not hold if the sequence of centers (c_n) converges to a projective measured lamination, see Corollary 7.3.

Fix a hyperbolic metric h on S and let c' be a point in the boundary of the augmented Teichmüller space of S . Then c' can be considered as a complete hyperbolic metric of finite area on $S \setminus \Gamma$, where Γ is the disjoint union of simple closed curves $\gamma_1, \dots, \gamma_l$. Up to isotopy, we can assume that γ_i is a geodesic for h and we denote by ℓ_i the h -length of γ_i . We will also denote by \bar{S} the surface obtained from S by collapsing each γ_i to a node ν_i .

We recall a construction of an infinite energy harmonic map $f : (S \setminus \Gamma, c') \rightarrow (S \setminus \Gamma, h)$ by Wolf [34]. For every $i = 1, \dots, l$, choose a sequence $s_{i,n} \rightarrow +\infty$ with $s_{i,n} > 1$. Let $U_{i,+}(s)$ and $U_{i,-}(s)$ be the cusps of $(S \setminus \Gamma, c')$ bounded by horocycles of length $1/s$ near ν_i and let $U_i(s) := U_{i,+}(s) \cup U_{i,-}(s)$ and $\bar{U}_i(s) := U_i(s) \cup \gamma_i$.

Fix an isometry $\xi_{i,\bullet} : U_{i,\bullet}(1) \rightarrow U/(0, y) \sim (1, y)$ with $U = [0, 1] \times [1, +\infty) \subset \mathbb{H}$ and put on $U_{i,\bullet}(1)$ the flat metric $|\Psi_i|$ induced by the quadratic differential $\Psi_i = \xi_{i,\bullet}^*(dz^2)$.

Define a new metric c on $S \setminus \Gamma$ which

- agrees with c' outside $\bigcup_i U_i(1)$ and with $|\Psi_i|$ on each $U_{i,\bullet}(1)$,
- is conformally equivalent to c' ,
- c is smooth away from $\partial U_i(1)$.

Let $U_{i,\bullet}^n(s)$ be the annulus $U_{i,\bullet}(s) \setminus U_{i,\bullet}(s_{i,n})$ and call $U_i^n(s) := U_{i,+}^n(s) \cup U_{i,-}^n(s)$. We denote by $\xi_{i,\bullet}^n : U_{i,\bullet}^n(1) \rightarrow U/\sim$ the restriction of $\xi_{i,\bullet}$.

Then S_n is obtained from S by removing $\bar{U}_i(s_{i,n})$ from the cusps adjacent to γ_i . Gluing the seams together, we obtain a compact surface \bar{S}_n with quadratic differentials Ψ_i^n on $\bar{U}_i^n(1)$ obtained restricting Ψ_i , with distinguished geodesics γ_i^n corresponding to the seams and collars $\bar{U}_i^n(s) = U_i^n(s) \cup \gamma_i^n$. We will also define $\xi_i^n : \bar{U}_i^n(1) \rightarrow U/\sim$ as

$$\xi_i^n(p) = \begin{cases} \xi_{i,+}^n(p) & \text{if } p \in \bar{U}_{i,+}^n \\ 1 + 2i(s_{i,n} + 1) - \xi_{i,-}^n(p) & \text{if } p \in \bar{U}_{i,-}^n \end{cases}$$

Notice that the metric c_n induced by c on \bar{S}_n determines a point in $\mathcal{T}(S)$: we will denote by c'_n the hyperbolic metric conformally equivalent to c_n .

Notice that, hidden in this construction, there is an arbitrary choice of twists associated to the gluings or, equivalently, to the charts $\xi_{i,\bullet}$.

Call $f_n : (\bar{S}_n, c_n) \rightarrow (S, h)$ the unique harmonic map in the given homotopy class [9].

Theorem 7.1 (Wolf [34]). *Up to subsequences, f_n converges $C^{2,\alpha}$ to a harmonic map $f : (S \setminus \Gamma, c) \rightarrow (S \setminus \Gamma, h)$ on the compact subsets of $S \setminus \Gamma$.*

In each cusp $U_{i,\bullet}(1)$ of $(S \setminus \Gamma, c)$, the Hopf differential Φ of f looks like $\Phi = (\ell_i^2/4 + O(e^{-2\pi y \circ \xi_{i,\bullet}}))\Psi_i$. Moreover, the energy density $e(f; c, h) \rightarrow \ell_i^2/2$ and the holomorphic energy density $H(f; c, h) \rightarrow \ell_i^2/4$ as $y \circ \xi_{i,\bullet} \rightarrow +\infty$.

We will assume throughout this section that we have already extracted a good subsequence (which we will still call f_n) so that the above theorem holds.

Now let h_n^* be the metric on S antipodal to h with respect to $(f_n)_*c_n$. Because of the theorem, h_n^* converges to some h^* smoothly away from Γ .

Proposition 7.2. *Let $1 = b_1 = \dots = b_r > b_{r+1} \geq \dots \geq b_l > 0$ and put $s_{i,n} = (a_i/\ell_i)t_n^{b_i}$, where $a_i > 0$ and $t_n \rightarrow +\infty$. Then, up to subsequences, $c_n \rightarrow [b_1\gamma_1 + \dots + b_l\gamma_l]$ and $h_n^* \rightarrow [a_1\gamma_1 + \dots + a_r\gamma_r]$ in Thurston compactification of $\mathcal{T}(S)$.*

Corollary 7.3. *If h is fixed and $h_n^* \rightarrow [\lambda]$ in Thurston compactification, then c_n does not necessarily converge to $[\lambda]$. If h is fixed and $c_n \rightarrow [\lambda]$ in Thurston compactification, then h_n^* does not necessarily converge to $[\lambda]$ and so the cyclic flow centered at c_n does not necessarily converge to an earthquake along λ (with any normalization).*

In order to prove Proposition 7.2, we need to estimate the transversal length $\text{trl}_{\gamma_i}(c'_n)$, that is, the width of a standard c'_n -collar of γ_i^n bounded by hypercycles of length 1.

Lemma 7.4. *The extremal length of γ_i at c_n satisfies*

$$\frac{1}{C_1 + 2(a_i/\ell_i)t_n^{b_i}} \leq \text{Ext}_{\gamma_i}(c_n) \leq \frac{\ell_i}{2a_it_n^{b_i}}$$

and so $\text{trl}_{\gamma_i}(c'_n) \asymp 2b_i \log t_n$.

Proof. By construction, $(\overline{U}_i^n(1), c_n)$ contains a flat cylinder of circumference 1 and height $2(a_i/\ell_i)t_n^{b_i}$ and so the extremal length satisfies

$$\text{Ext}_{\gamma_i}(c_n) \leq \frac{\ell_i}{2a_i t_n^{b_i}}.$$

On the other hand, consider a metric \hat{c}_n on \overline{S}_n which is conformally equivalent to c_n , which agrees with c_n on $\overline{S}_n \setminus \bigcup_{j \neq i} \overline{U}_j^n$ and such that the \hat{c}_n -area of \overline{U}_j^n is bounded by a fixed constant for $j \neq i$ and the distance between the two boundary components of \overline{U}_j^n is at least 1. For instance, one can define \hat{c}_n be rescaling c_n by a factor which is constantly $1/s_{j,n}$ on the regions $\overline{U}_j^n(2)$ for all $j \neq i$, which interpolates between 1 and $1/s_{j,n}$ on $\overline{U}_j^n(1) \setminus \overline{U}_j^n(2)$ for $j \neq i$, and which is constantly 1 elsewhere.

Then $\ell_{\gamma_i}(\hat{c}_n) = 1$ and the area $\text{Area}_{\hat{c}_n}(S) \leq C_1 + 2(a_i/\ell_i)t_n^{b_i}$ where C_1 is a constant that depends only on $\chi(S)$ and k . Hence,

$$\text{Ext}_{\gamma_i}(c_n) \geq \frac{1}{C_1 + 2(a_i/\ell_i)t_n^{b_i}}$$

As $\text{Ext}_{\gamma_i}(c_n) \rightarrow 0$, Maskit's estimate (see [18]) gives $\ell_{\gamma_i}(c_n) \asymp \pi \text{Ext}_{\gamma_i}(c_n)$ and so $\text{trl}_{\gamma_i}(c'_n) \asymp -2 \log \ell_{\gamma_i}(c'_n) \asymp 2b_i \log t_n$. \square

For each i , fix an open neighbourhood $A_i \subset (\overline{S}, c)$ of ν_i whose closure does not contain any zero of Φ , and such that $(A_i \setminus \nu_i, |\Phi|)$ is the union of two annuli. Moreover, choose standard h -collars $R_i \subset (S, h)$ around γ_i such that $R_i \subset f(A_i)$. By Wolf's construction (see [34]), outside $\bigcup_i f^{-1}(R_i)$ the Hopf differential Φ_n of f_n converges $C^{1,\alpha}$ to Φ .

Here we recall that, by definition,

$$h = 2(f_n)_* \text{Re}(\Phi_n) + e(f_n; c_n, h) c_n$$

where $e(f_n; c_n, h)$ is the energy density; moreover, by Equation (2),

$$(f_n)_* \text{Re}(\Phi_n) = \frac{1}{4} h((E - b_n^2) \bullet, \bullet)$$

and so b_n converges $C^{1,\alpha}$ to b outside $\bigcup_i R_i$.

Notice that the horizontal (resp. vertical) directions of Φ_n are exactly the eigenspaces of b_n corresponding to the smaller (resp. bigger) eigenvalue.

Lemma 7.5. *Fix a small $\varepsilon > 0$. Up to shrinking R_i and for n large enough,*

$$\left| \frac{4}{\ell_i^2} \frac{\Phi_n}{\Psi_i^n} - 1 \right| < \varepsilon^2$$

and

$$\left| \frac{2}{\ell_i^2} e(f_n; c_n, h) - 1 \right| < 2\varepsilon^2$$

in every $f_n^{-1}(R_i)$.

Proof. Up to shrinking R_i , we can assume that $R_i \subset f_n(\overline{U}_i^n(3))$ and there

$$\left| \frac{4}{\ell_i^2} \frac{\Phi}{\Psi_i} - 1 \right| < \varepsilon^2/2 \quad \text{and} \quad \left| \frac{4}{\ell_i^2} H(f; c, h) - 1 \right| < \varepsilon^2/2$$

where $H(f; c, h) = \frac{1}{2} \|\partial f\|^2$ is the holomorphic energy density of f .

As $f_n^{-1}|\partial R_i \rightarrow f^{-1}|\partial R_i$, for n large enough $f_n^{-1}(\partial R_i)$ is contained inside $\overline{U}_i^n(2)$. Moreover, $(\xi_{i,\bullet}^n)_* \Phi_n \rightarrow (\xi_{i,\bullet})_* \Phi$ in a compact neighbourhood $K_{i,\bullet}$ of $\xi_{i,\bullet}^n(f_n^{-1}(\partial R_i) \cap U_{i,\bullet}^n)$ and so

$$\frac{4}{\ell_i^2} \left| (\xi_{i,\bullet}^n)_* \frac{\Phi_n}{\Psi_i^n} - (\xi_{i,\bullet})_* \frac{\Phi}{\Psi_i} \right| < \varepsilon^2/2$$

in $K_{i,\bullet}$. Thus,

$$\frac{4}{\ell_i^2} \left| (\xi_i^n)_* \frac{\Phi_n}{\Psi_i^n} - 1 \right| < \varepsilon^2$$

in $K_{i,\bullet}$. Because $\xi_{i,\bullet}^n$ is holomorphic and Φ_n/Ψ_i^n is a holomorphic function on $f_n^{-1}(R_i)$, the same estimate holds in $f_n^{-1}(R_i)$ for n large enough.

In a similar way, if $H_n = H(f_n; c_n, h)$, then the energy density and the Jacobian of f_n are

$$e(f_n; c_n, h) = H_n + \frac{|\Phi_n|^2}{|\Psi_i^n|^2 H_n} \quad \text{and} \quad J(f_n; c_n, h) = H_n - \frac{|\Phi_n|^2}{|\Psi_i^n|^2 H_n} > 0$$

which implies $H_n > |\Phi_n|/|\Psi_i^n| > (\ell_i^2/4)(1 - \varepsilon^2)$ on $f_n^{-1}(R_i)$.

On the other hand (see [33], for instance)

$$\Delta_{c_n} \log H_n = 2J_n > 0$$

and so $\Delta_{c_n} \log(4H_n/\ell_i^2) \geq 0$ on $\overline{U}_i^n(2)$. As $\log(4H_n/\ell_i^2) < \varepsilon^2/2$ on $\partial f_n^{-1}(R_i)$ for n large, we obtain $H_n \leq (\ell_i^2/4)(1 + \varepsilon^2/2)$ on $f_n^{-1}(R_i)$ and so the wished estimate for $e(f_n; c_n, h)$. \square

Thanks to the previous lemma, we can draw a few consequences.

Corollary 7.6. (i) *The metric h^* has nodes at Γ and $h_n^* \rightarrow h^*$ in the augmented Teichmüller space.*
(ii) *In the whole $f_n^{-1}(R_i)$ found in the above lemma, the bigger eigenvalue κ of b_n is greater than $1/2\varepsilon$ for large n .*

Proof. As for (i), notice that $h_n^* = -2\text{Re}(\Phi_n) + e(f_n; c_n, h)c_n$. Because of Lemma 7.5, for every $\varepsilon > 0$ there exists R_i such that, for $n > n(\varepsilon)$, the h_n^* -norm of $(f_n \circ (\xi_i^n)^{-1})_* \partial_x$ is at most $2\varepsilon\ell_i$ the h_n -norm of $(f_n \circ (\xi_i^n)^{-1})_* \partial_y$ is at most $2\varepsilon\ell_i$. Thus, $\ell_{\gamma_i}(h_n^*) \leq 2\varepsilon\ell_i$ and so $\ell_{\gamma_i}(h_n^*) \rightarrow 0$. As $h_n^* \rightarrow h^*$ on the compact subsets of $S \setminus \Gamma$, we conclude that h^* has nodes at γ_i and the sequence converges in the augmented Teichmüller space.

As for (ii), the two identities $f_n^* h = 2\text{Re}(\Phi_n) + e(f_n; c_n, h)c_n$ and $f_n^* h((E - b_n^2)_\bullet, \bullet) = 4\text{Re}(\Phi_n)$, already seen in Section 3 Equation (2), give

$$\frac{h((E - b_n^2)(f_n)_* v, (f_n)_* v)}{h((f_n)_* v, (f_n)_* v)} = \frac{4\text{Re}(\Phi_n)(v, v)}{2\text{Re}(\Phi_n)(v, v) + e(f_n; c_n, h)c_n(v, v)}$$

for any vector tangent v to $\overline{U}_i^n(1)$. Choosing $v = (\xi_i^n)^* \partial_y$, we obtain

$$\frac{h((E - b_n^2)(f_n)_* v, (f_n)_* v)}{h((f_n)_* v, (f_n)_* v)} \leq \frac{\ell_i^2(-1 + 3\varepsilon^2)}{4\ell_i^2\varepsilon^2}$$

on $\overline{U}_i^n(1) \cap f_n^{-1}(R_i)$ and so $1 - \kappa^2 \leq (-1/4\varepsilon^2) + 1$ there, that is $\kappa \geq 1/2\varepsilon$. \square

Up to subsequences, we can assume that h_n^* converges to a point $[\lambda]$ in Thurston boundary. Notice that λ must be supported on $\gamma_1 \cup \dots \cup \gamma_l$ and so $\lambda = w_1\gamma_1 + \dots + w_l\gamma_l$ with $w_1, \dots, w_l \geq 0$.

We will show that $[\lambda] = [a_1\gamma_1 + \dots + a_r\gamma_r]$, and so the result will not depend on the chosen subsequence.

Let $(\theta_n)_{n \in \mathbb{N}}$ be a sequence of positive numbers such that $\theta_n \ell_{h_n^*} \rightarrow \iota(\lambda, \bullet)$.

Lemma 7.7. *For every i , we have $\theta_n \ell_i s_{i,n} \rightarrow w_i$ as $n \rightarrow \infty$.*

Proof. Let $\varepsilon > 0$. We can choose a collar $\gamma_i \subset R_i \subset S$ such that $\text{tr}(b_n) > 2/\varepsilon$ on R_i and $|\frac{4}{\ell_i^2}\Phi_n - \Psi_i^n| < \varepsilon^2|\Psi_i^n|$ on $f_n^{-1}(R_i)$ for n large.

As $f_n^{-1} \rightarrow f^{-1}$ on $S \setminus \Gamma$, we can assume that there exists $\bar{y} > 1$ such that $\overline{U}_n^i(1) \supset f_n^{-1}(R_i) \supset \overline{U}_n^i(\bar{y})$ for n large.

By abuse of notation, denote just by (x, y) the Euclidean coordinates on R_i given by the parametrization $\xi_i^n \circ f_n$.

By writing the relation $4\text{Re}(\Phi_n) = h((E - b_n^2)_\bullet, \bullet)$ in coordinates on R_i and taking the determinant, we obtain

$$(52) \quad (1 + \eta(\varepsilon))\ell_i^2 dx \wedge dy = \sqrt{\text{tr}(b_n)^2 - 4} \omega_h$$

where $|\eta(\varepsilon)| < \varepsilon^2$ for n large, by Lemma 7.5.

By Corollary 7.6(ii), $(1 - \varepsilon^2/2)\text{tr}(b_n) \leq \sqrt{\text{tr}(b_n)^2 - 4} \leq \text{tr}(b_n)$ for n large.

Multiplying by θ_n both hand sides of (52) and integrating over R_i , we get

$$(53) \quad \theta_n(1 - \varepsilon^2/2) \int_{R_i} \text{tr}(b_n)\omega_h \leq \theta_n(1 + \eta(\varepsilon))\ell_i^2 \int_{R_i} dx \wedge dy \leq \theta_n \int_{R_i} \text{tr}(b_n)\omega_h$$

Now,

$$(54) \quad \theta_n \ell_i^2(s_{i,n} - \bar{y}) \leq \theta_n \ell_i^2 \int_{R_i} dx \wedge dy \leq \theta_n \ell_i^2 s_{i,n}$$

and $\theta_n \int_{R_i} \text{tr}(b_n)\omega_h \rightarrow w_i \ell_i$ by Proposition 6.15.

The result follows by comparing Equations (53) and (54). \square

Proof of Proposition 7.2. By Lemma 7.4 and Maskit's estimate, we immediately obtain that $c_n \rightarrow [b_1\gamma_1 + \dots + b_l\gamma_l]$.

By Lemma 7.7,

$$\frac{w_i}{w_j} = \lim_n \frac{\ell_i s_{i,n}}{\ell_j s_{j,n}} = \lim_n \frac{a_i t_n^{b_i}}{a_j t_n^{b_j}}$$

which shows that $[\lambda] = [a_1\gamma_1 + \dots + a_r\gamma_r]$. \square

8. THE LANDSLIDE FLOW ON THE UNIVERSAL TEICHMÜLLER SPACE

In this section we show how the construction of the landslide flow L extends to the universal Teichmüller space. We believe that this S^1 action on the product of two copies of the universal Teichmüller space can be of independent interest, but limit our investigations here to its definition and to checking that it is non-trivial.

8.1. Minimal Lagrangian maps and the universal Teichmüller space. The universal Teichmüller space \mathcal{T}_U can be defined as the quotient of the group \mathcal{QS} of quasi-symmetric homeomorphisms of the circle by composition on the left by projective transformations, see e.g. [10]. We will show here that the map L defined above extends to a circle action \mathcal{L} on $\mathcal{T}_U \times \mathcal{T}_U$. This is based on the following statement.

Theorem 8.1 ([7]). *Let $\bar{\psi} \in \mathcal{QS}$. There exists a unique quasi-conformal minimal Lagrangian diffeomorphism $m : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ such that $\partial m = \bar{\psi}$.*

We call g the hyperbolic metric on \mathbb{H}^2 , and ∇ its Levi-Civita connection. It follows from the basic facts on minimal Lagrangian diffeomorphisms, as recalled in Section 1.3, that there exists a unique bundle morphism $b : T\mathbb{H}^2 \rightarrow T\mathbb{H}^2$ such that:

- $m^*g = g(b\bullet, b\bullet)$,
- $\det(b) = 1$,
- b is self-adjoint for g ,
- b satisfies the Codazzi equation: $d^\nabla b = 0$.

Since m is quasi-conformal, b has eigenvalues in $[\epsilon, 1/\epsilon]$ for some $\epsilon > 0$, see [7].

8.2. An extension of L to the universal Teichmüller space. The construction of the S^1 -action is based on the following definition and lemma.

Definition 8.2. Let $\bar{\psi} \in \mathcal{QS}$ and let $e^{i\theta} \in S^1$. Let $m : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ be the unique quasi-conformal minimal Lagrangian diffeomorphism such that $\partial m = \bar{\psi}$, and let $b : T\mathbb{H}^2 \rightarrow T\mathbb{H}^2$ be as above. We call

$$\beta_\theta = \cos(\theta/2)E + \sin(\theta/2)Jb$$

and set $g_\theta := g(\beta_\theta\bullet, \beta_\theta\bullet)$ and $g_\theta^* = g(\beta_{\theta+\pi}\bullet, \beta_{\theta+\pi}\bullet)$.

Lemma 8.3. *With the notations above, g_θ and g_θ^* are complete hyperbolic metrics on \mathbb{H}^2 . The identity map from (\mathbb{H}^2, g_θ) to $(\mathbb{H}^2, g_\theta^*)$ is minimal Lagrangian and quasi-conformal.*

Sketch of the proof. The fact that g_θ and g_θ^* have curvature -1 follows from the same argument as in the proof of Proposition 1.7; we do not repeat it here. Moreover, the argument given in the proof of Theorem 1.10 also shows that the identity map from (\mathbb{H}^2, g_θ) to $(\mathbb{H}^2, g_\theta^*)$ is minimal Lagrangian.

To check that the identity between (\mathbb{H}^2, g) and (\mathbb{H}^2, g_θ) is quasiconformal, it is sufficient to prove that the eigenvalues of the bundle morphism ${}^t\beta_\theta \cdot \beta_\theta$ are between ϵ' and $1/\epsilon'$, for some $\epsilon' > 0$ depending on ϵ . However $\det(\beta_\theta) = 1$ by definition, so that ${}^t\beta_\theta \cdot \beta_\theta$ has determinant 1. To compute its trace, notice that

$${}^t\beta_\theta \cdot \beta_\theta = \cos^2(\theta/2)E + \sin^2(\theta/2)b^2 + \cos(\theta/2)\sin(\theta/2)(Jb - bJ)$$

and that $\text{tr}(Jb) = \text{tr}(bJ) = 0$. It follows that

$$\text{tr}({}^t\beta_\theta \cdot \beta_\theta) = 2\cos^2(\theta/2) + \sin^2(\theta/2)\text{tr}(b^2).$$

Since the eigenvalues of b are in $[\epsilon, 1/\epsilon]$, it follows that the identity between (\mathbb{H}^2, g) and (\mathbb{H}^2, g_θ) is quasi-conformal, and that g_θ and g_θ^* are complete. \square

As a consequence we can give the definition of the action considered here. Let $\psi, \psi^* \in \mathcal{QS}$ represent points $[\psi], [\psi^*] \in \mathcal{T}_U$. Then $\bar{\psi} := \psi^* \circ \psi^{-1} : S^1 \rightarrow S^1$ is a quasimetric homeomorphism: let $m : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ be the unique quasiconformal minimal Lagrangian diffeomorphism with $\partial m = \bar{\psi}$. We can then define a bundle morphism $b : T\mathbb{H}^2 \rightarrow T\mathbb{H}^2$ as above, as well as two hyperbolic metrics g_θ and g_θ^* associated to θ and b , as in Section 3. Lemma 8.3 shows that the identity between (\mathbb{H}^2, g_θ) and $(\mathbb{H}^2, g_\theta^*)$ is minimal Lagrangian.

Since g_θ is hyperbolic, (\mathbb{H}^2, g_θ) is isometric to the hyperbolic plane. The identity between (\mathbb{H}^2, g) and (\mathbb{H}^2, g_θ) therefore determines a quasiconformal diffeomorphism Ψ_θ between (\mathbb{H}^2, g) and \mathbb{H}^2 , well-defined up to composition on the left, with boundary value $\psi_\theta \in \mathcal{QS}$. Similarly, the identity map from (\mathbb{H}^2, g) to $(\mathbb{H}^2, g_\theta^*)$ determines a quasiconformal map Ψ_θ^* between the hyperbolic plane and itself, with boundary value $\psi_\theta^* \in \mathcal{QS}$. Then $\Psi_\theta^* \circ \Psi_\theta^{-1}$ is a quasiconformal minimal Lagrangian diffeomorphism by Lemma 8.3. We define \mathcal{L} as:

$$\mathcal{L}_{e^{i\theta}}([\psi], [\psi^*]) = ([\psi_\theta], [\psi_\theta^*]) .$$

To establish a relation between \mathcal{T}_U and \mathcal{T}_S , fix a hyperbolic metric h_0 on S and a universal covering map $\mathbb{H}^2 \cong (\tilde{S}, \tilde{h}_0) \rightarrow (S, h_0)$, so that $\rho_0 : \pi_1(S) \rightarrow \mathrm{PSL}_2(\mathbb{R})$ is the associated holonomy representation. Given $[h] \in \mathcal{T}_S$, we can consider the lift \tilde{h} of h to \mathbb{H}^2 and let ρ be its holonomy representation. The identity map from $(\mathbb{H}^2, \tilde{h}_0)$ to $(\mathbb{H}^2, \tilde{h})$ determines a quasi-conformal diffeomorphism Ψ of \mathbb{H}^2 to itself, with boundary value $\psi \in \mathcal{QS}$, that conjugates the action of ρ_0 on \mathbb{H}^2 to the action of ρ .

Let $i_S : \mathcal{T}_S \hookrightarrow \mathcal{T}_U$ be the canonical embedding of Teichmüller space of S in the universal Teichmüller space defined as $i_S([h]) = [\psi]$.

Proposition 8.4. *The restriction via i_S of \mathcal{L} to $\mathcal{T}_S \times \mathcal{T}_S \subset \mathcal{T}_U \times \mathcal{T}_U$ is the landslide action L .*

Proof. Let $[\psi] = i_S([h])$ and $[\psi^*] = i_S([h^*])$ be points of $i_S(\mathcal{T}_S) \subset \mathcal{T}_U$. Let ρ and ρ^* be the holonomy representations of h and h^* respectively. Then ψ and ψ^* are the boundary values of quasiconformal maps $\Psi, \Psi^* : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ which are conjugating ρ_0 to actions ρ and ρ^* on \mathbb{H}^2 . By construction, $g_\theta = \tilde{h}_\theta$ is ρ -invariant and $g_\theta^* = \tilde{h}_\theta^*$ is ρ^* -invariant; moreover, Ψ_θ (resp. Ψ_θ^*) conjugates the action of ρ_0 to the action of the holonomy representation ρ_θ of h_θ (resp. the holonomy representation ρ_θ^* of h_θ^*). Hence, $([\psi_\theta], [\psi_\theta^*]) \in i_S(\mathcal{T}_S) \times i_S(\mathcal{T}_S) \subset \mathcal{T}_U \times \mathcal{T}_U$, and the restriction of \mathcal{L} to $\mathcal{T}_S \times \mathcal{T}_S$ coincides with L , as claimed. \square

Theorem 8.5. *The map \mathcal{L} defines a non-trivial action of S^1 on \mathcal{T}_U .*

Sketch of the proof. To prove that \mathcal{L} determines an action of S^1 , it is sufficient to check that, for all $\theta, \theta' \in \mathbb{R}$ and all $[\psi], [\psi^*] \in \mathcal{T}_U$, $\mathcal{L}_{e^{i\theta}}(\mathcal{L}_{e^{i\theta'}}([\psi], [\psi^*])) = \mathcal{L}_{e^{i(\theta+\theta')}}([\psi], [\psi^*])$.

However, this follows from the fact that $\Psi_\theta^* \circ \Psi_\theta^{-1}$ is minimal Lagrangian, followed by the same argument used in the proof of Theorem 1.8; so we do not repeat them here.

The nontriviality of \mathcal{L} is clear, since L is non-trivial in all the copies of Teichmüller spaces of surfaces of finite genus, see Proposition 8.4. \square

9. APPLICATIONS, EXTENSIONS, AND QUESTIONS

This section contains a brief outline of some possible applications of the landslide flow developed here, and of some open questions.

9.1. Constant curvature surfaces in globally hyperbolic AdS manifolds. As already mentioned, a recent result of Barbot, Béguin and Zeghib [3] states that the complement of the convex core in a globally hyperbolic AdS manifold has a unique foliation by constant Gauss curvature surfaces. This was used above in the proof of Theorem 1.14.

However we believe that this result might not be necessary to prove Theorem 1.14, and that a direct, albeit longer, proof could be given, based on a deformation argument. Using the same argument as in the proof of Theorem 1.14, but backwards, it should then be possible to recover parts of the main result of [3]: the proof of the existence, in a given globally hyperbolic AdS 3-dimensional manifold, of a unique surface of prescribed constant curvature.

9.2. Holomorphic disks in Teichmüller space. One obvious consequence of Theorems 1.14 and 5.1 is the existence of many holomorphic disks in Teichmüller space of S : given $h, h' \in \mathcal{T}$ with $h \neq h'$ and given $\zeta \in S^1 \setminus \{1\}$ there exists $h^* \in \mathcal{T}$ and a holomorphic map $C_\bullet(h, h^*) : \overline{\Delta} \rightarrow \mathcal{T}$ from the unit disk in \mathbb{C} to Teichmüller space of S , such that $C_1(h, h^*) = h$ and $C_\zeta(h, h^*) = h'$.

It is of course conceivable that the disks obtained in this manner for two different values of h' have the same image. However, there are reasons to believe that it is not often the case. If this is correct, it would mean that the landslide disks provide a $(12g - 11)$ -dimensional family of holomorphic disks in \mathcal{T} .

9.3. Other questions. There are many remaining questions concerning the landslide flow or its complex extension, mostly motivated by the analogy with the earthquake flow. Some of those statements can be translated in terms of 3-dimensional hyperbolic or AdS geometry. We give here a short list of example of possible questions.

Smooth grafting as homeomorphism. Recall that Scannell and Wolf [26] proved that, for $\lambda \in \mathcal{ML}$ fixed, the map $h \mapsto gr_\lambda(h)$ is a homeomorphism of \mathcal{T} . When $h \in \mathcal{T}$ is fixed, the map $\lambda \mapsto gr_\lambda(h)$ is also a homeomorphism from \mathcal{ML} to \mathcal{T} , see [8].

It is tempting to ask whether those statements can be extended to the smooth grafting map sgr . Note that in this setting the two statements above concerning the grafting map – with the measured lamination fixed, and with the hyperbolic metric fixed – are now merged into one, since the two hyperbolic metrics that occur in the map sgr play symmetric roles.

Question 9.1. Let $s \in (0, 1)$, and let $h \in \mathcal{T}$. Is the map $h^* \mapsto sgr_s(h, h^*)$ a homeomorphism from \mathcal{T} to \mathcal{T} ?

This statement can be translated in terms of the geometry of hyperbolic ends, in the following, essentially equivalent question.

Question 9.2. Let $h, c \in \mathcal{T}$ and let $K \in (-1, 0)$.

- Is there a unique hyperbolic end with conformal structure at infinity c , and containing an embedded surface of constant curvature K with induced metric proportional to h ?
- Is there a unique hyperbolic end with conformal structure at infinity c , containing an embedded surface of constant curvature K with third fundamental form proportional to h^* ?

The action of the landslide flow at infinity. It is quite natural to wonder to what extent the landslide flow can be extended to Thurston boundary of Teichmüller space. One side of this question is already answered above in Section 6, concerning the limit of L to the earthquake flow when one of the parameter converges to Thurston boundary and the other is fixed. However other questions can be asked, in particular in light of the results of Wolf [34] on the behavior of harmonic maps at the boundary of \mathcal{T} .

The landslide flow as a Hamiltonian flow. Consider a fixed measured lamination $\lambda \in \mathcal{ML}$. The flow of earthquakes along λ is the Hamiltonian flow of the length of λ , considered as a function on \mathcal{T} , with respect to the Weil-Petersson symplectic structure. In a similar way, is the landslide flow the Hamiltonian flow of some functional on $\mathcal{T} \times \mathcal{T}$?

The data at infinity of hyperbolic ends. For all $K \in (-1, 0)$, there is a parameterization of \mathcal{CP} by $\mathcal{T} \times \mathcal{T}$, with a complex projective structure P corresponding to (h, h^*) if the hyperbolic end E with complex projective structure P at infinity contains a surface of constant curvature K with induced metric proportional to h and third fundamental form proportional to h^* .

There is also another parameterization of \mathcal{CP} by the space of couples (h, b) , where $h \in \mathcal{T}$ and where b is a bundle morphism which is self-adjoint for h and satisfies the Codazzi equation and $\det(b) = 1$.

Given a \mathbb{CP}^1 -structure P , we can also consider the data at infinity I^* and III^* of the corresponding hyperbolic end, as defined in [13], and take the limit as $K \rightarrow 0$. Is it true that h and h^* limit to I^* and III^* respectively? And that the traceless part of B , suitably renormalized, converges to B^* ?

Landslides on the universal Teichmüller space. Section 8 on the universal Teichmüller space leaves a number of elementary questions unanswered. One natural question is whether for fixed $[\psi], [\psi^*] \in \mathcal{T}_U$ the map $e^{i\theta} \rightarrow \mathcal{L}_{e^{i\theta}}^1([\psi], [\psi^*])$ extends to a holomorphic disk in \mathcal{T}_U , as for the landslide action on the Teichmüller space of a closed surface.

Another natural question is whether all fixed points of the landslide action on $\mathcal{T}_U \times \mathcal{T}_U$ are on the diagonal.

Cone singularities. It appears possible that all the results obtained here extend from closed hyperbolic surfaces to finite volume hyperbolic surfaces, and more generally to hyperbolic surfaces with cone singularities (perhaps of angle less than π). The 3-dimensional AdS or hyperbolic part of the picture would then be filled with 3-dimensional AdS or hyperbolic manifolds with “particles”, as considered e.g. in [12, 22, 17].

REFERENCES

1. Lars Andersson, Thierry Barbot, Riccardo Benedetti, Francesco Bonsante, William M. Goldman, François Labourie, Kevin P. Scannell, and Jean-Marc Schlenker, *Notes on: “Lorentz spacetimes of constant curvature”* [Geom. Dedicata **126** (2007), 3–45; mr2328921] by G. Mess, Geom. Dedicata **126** (2007), 47–70. MR MR2328922
2. Thierry Barbot, François Béguin, and Abdelghani Zeghib, *Constant mean curvature foliations of globally hyperbolic spacetimes locally modelled on AdS_3* , Geom. Dedicata **126** (2007), 71–129. MR MR2328923 (2008j:53041)
3. Thierry Barbot, François Béguin, and Abdelghani Zeghib, *Prescribing Gauss curvature of surfaces in 3-dimensional spacetimes, application to the Minkowski problem in the Minkowski space*, 2008, arXiv.org:0804.1053. To appear, Ann. Institut Fourier.
4. Mehdi Belraouti, PhD thesis, in preparation.
5. Riccardo Benedetti and Francesco Bonsante, *Canonical Wick rotations in 3-dimensional gravity*, Memoirs of the American Mathematical Society **198** (2009), 164pp, math.DG/0508485.

6. Riccardo Benedetti and Enore Guadagnini, *Cosmological time in $(2+1)$ -gravity*, Nuclear Phys. B **613** (2001), no. 1-2, 330–352. MR MR1857817 (2002i:83047)
7. Francesco Bonsante and Jean-Marc Schlenker, *Maximal surfaces and the universal Teichmüller space*, Invent. Math. **182** (2010), no. 2, 279–333. MR 2729269
8. David Dumas and Michael Wolf, *Projective structures, grafting and measured laminations*, Geom. Topol. **12** (2008), no. 1, 351–386. MR MR2390348 (2009c:30114)
9. James Eells, Jr. and J. H. Sampson, *Harmonic mappings of Riemannian manifolds*, Amer. J. Math. **86** (1964), 109–160. MR 0164306 (29 #1603)
10. Frederick P. Gardiner and William J. Harvey, *Universal Teichmüller space*, Handbook of complex analysis: geometric function theory, Vol. 1, North-Holland, Amsterdam, 2002, pp. 457–492. MR 1966201 (2004a:30041)
11. Steven P. Kerckhoff, *The Nielsen realization problem*, Ann. of Math. (2) **117** (1983), no. 2, 235–265. MR MR690845 (85e:32029)
12. Kirill Krasnov and Jean-Marc Schlenker, *Minimal surfaces and particles in 3-manifolds*, Geom. Dedicata **126** (2007), 187–254. MR MR2328927
13. ———, *On the renormalized volume of hyperbolic 3-manifolds*, Comm. Math. Phys. **279** (2008), no. 3, 637–668. MR MR2386723
14. Ravi S. Kulkarni and Ulrich Pinkall, *A canonical metric for Möbius structures and its applications*, Math. Z. **216** (1994), no. 1, 89–129.
15. F. Labourie, *Problème de Minkowski et surfaces à courbure constante dans les variétés hyperboliques*, Bull. Soc. Math. Fr. **119**(1991), 307–325
16. F. Labourie, *Surfaces convexes dans l'espace hyperbolique et CP^1 -structures*, J. London Math. Soc., II. Ser. **45** (1992), 549–565.
17. Cyril Lecuire and Jean-Marc Schlenker, *The convex core of quasifuchsian manifolds with particles*, arXiv:0909.4182, 2009.
18. Bernard Maskit, *Comparison of hyperbolic and extremal lengths*, Ann. Acad. Sci. Fenn. Ser. A I Math. **10** (1985), 381–386. MR 87c:30062
19. Curtis T. McMullen, *Complex earthquakes and Teichmüller theory*, J. Amer. Math. Soc. **11** (1998), no. 2, 283–320. MR 1478844 (98i:32030)
20. Geoffrey Mess, *Lorentz spacetimes of constant curvature*, Geom. Dedicata **126** (2007), 3–45. MR MR2328921
21. Gabriele Mondello, *Flows of $SL_2(\mathbb{R})$ -type on the cotangent space to Teichmüller space*, in preparation.
22. Sergiu Moroianu and Jean-Marc Schlenker, *Quasi-Fuchsian manifolds with particles*, J. Differential Geom. **83** (2009), no. 1, 75–129. MR MR2545031
23. Igor Rivin and Craig D. Hodgson, *A characterization of compact convex polyhedra in hyperbolic 3-space*, Invent. Math. **111** (1993), 77–111.
24. J. H. Sampson, *Some properties and applications of harmonic mappings*, Ann. Sci. École Norm. Sup. (4) **11** (1978), no. 2, 211–228. MR 510549 (80b:58031)
25. Kevin P. Scannell, *Flat conformal structures and the classification of de Sitter manifolds*, Comm. Anal. Geom. **7** (1999), no. 2, 325–345. MR 1685590 (2000a:53122)
26. Kevin P. Scannell and Michael Wolf, *The grafting map of Teichmüller space*, J. Amer. Math. Soc. **15** (2002), no. 4, 893–927 (electronic). MR MR1915822 (2003d:32011)
27. Jean-Marc Schlenker, *Métriques sur les polyèdres hyperboliques convexes*, J. Differential Geom. **48** (1998), no. 2, 323–405. MR MR1630178 (2000a:52018)
28. ———, *Hypersurfaces in H^n and the space of its horospheres*, Geom. Funct. Anal. **12** (2002), no. 2, 395–435. MR MR1911666 (2003d:53108)
29. Richard Schoen and Shing Tung Yau, *On univalent harmonic maps between surfaces*, Invent. Math. **44** (1978), no. 3, 265–278. MR 0478219 (57 #17706)
30. Richard M. Schoen, *The role of harmonic mappings in rigidity and deformation problems*, Complex geometry (Osaka, 1990), Lecture Notes in Pure and Appl. Math., vol. 143, Dekker, New York, 1993, pp. 179–200. MR MR1201611 (94g:58055)
31. M. Spivak, *A comprehensive introduction to geometry, vol. i-v*, Publish or perish, 1970-1975.
32. William P. Thurston, *Three-dimensional geometry and topology.*, Originally notes of lectures at Princeton University, 1979. Recent version available on <http://www.msri.org/publications/books/gt3m/>, 1980.
33. Michael Wolf, *The Teichmüller theory of harmonic maps*, J. Differential Geom. **29** (1989), no. 2, 449–479. MR 982185 (90h:58023)
34. ———, *Infinite energy harmonic maps and degeneration of hyperbolic surfaces in moduli space*, J. Differential Geom. **33** (1991), no. 2, 487–539. MR 1094467 (92b:58055)

UNIVERSITÀ DEGLI STUDI DI PAVIA, VIA FERRATA, 1, 27100 PAVIA, ITALY

E-mail address: bonsante@sns.it

UNIVERSITÀ DI ROMA “LA SAPIENZA” - DIPARTIMENTO DI MATEMATICA “GUIDO CASTELNUOVO”, PIAZZALE ALDO MORO 5, 00185 ROMA, ITALY

E-mail address: mondello@mat.uniroma1.it

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR CNRS 5219, UNIVERSITÉ TOULOUSE III, 31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: schlenker@math.univ-toulouse.fr